



ORIGINAL ARTICLE

Sociodemographic, anthropometric and dietary determinants of dyslipidemia in preschoolers[☆]

Luciana N. Nobre^{a,*}, Joel A. Lamounier^b, Sylvia do C.C. Franceschini^c

^a Doctor. Departamento de Nutrição, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Diamantina, MG, Brazil

^b Doctor. Departamento de Medicina, Universidade Federal de São João Del-Rei, Divinópolis, MG, Brazil

^c Doctor. Departamento de Nutrição, Universidade Federal de Viçosa, Viçosa, MG, Brazil

Received 29 November 2012; accepted 6 February 2013

Available online 10 July 2013

KEYWORDS

Hypercholesterolemia;
Body mass index;
Feeding behavior

Abstract

Objective: To investigate the determinants of dyslipidemia in preschoolers.

Methods: A total of 227 preschoolers residing in an urban area of the city of Diamantina, Minas Gerais, Brazil were evaluated at age 5 years, using a cross-sectional design. Dietary intake from a food frequency questionnaire, anthropometric/biochemical parameters, and socioeconomic/behavioral information from a questionnaire were evaluated. 'Mixed diet', 'snack', and 'unhealthy' dietary patterns were identified using principal component analysis. The determinants of dyslipidemia were examined using Poisson regression analysis.

Results: The prevalence of dyslipidemia in this study was 65.19%. Preschoolers who less frequently consumed foods in the 'mixed diet' dietary pattern had a higher risk of high concentrations of low-density lipoprotein cholesterol (PR = 2.30; p = 0.004) when compared with those with more frequent consumption of the 'mixed diet' dietary pattern. Preschoolers whose mothers had lower levels of education presented a lower risk of high concentrations of low-density lipoprotein cholesterol (PR = 0.43; p = 0.003), and preschoolers who were overweight/obese presented with greater risk of high concentrations of low-density lipoprotein cholesterol (PR = 2.23; p = 0.003).

Conclusion: The determinants of dyslipidemia identified in this study were less frequent consumption of foods in the 'mixed diet' dietary pattern, higher body mass index, and lower level of maternal education. This study shows that despite the young age of the group under study, they already present a high prevalence of dyslipidemia, which is an important risk factor for cardiovascular disease.

© 2013 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

[☆] Please cite this article as: Nobre LN, Lamounier JA, Franceschini SC. Sociodemographic, anthropometric and dietary determinants of dyslipidemia in preschoolers. J Pediatr (Rio J). 2013;89:462–9.

* Corresponding author.

E-mail: lunerinobre@yahoo.com.br (L.N. Nobre).

PALAVRAS-CHAVE

Hipercolesterolemia;
Índice de massa
corporal;
Comportamento
alimentar

Determinantes sociodemográficos, antropométricos e alimentares de dislipidemia em pré-escolares**Resumo**

Objetivo: O objetivo deste estudo foi investigar os determinantes de dislipidemia em pré-escolares.

Métodos: A partir de um desenho transversal, foi avaliado um total de 227 pré-escolares com 5 anos de idade residentes em uma área urbana da cidade de Diamantina, Minas Gerais, Brasil. Foram avaliados: variáveis bioquímicas, antropométricas, condição socioeconômica e comportamental, a partir de um questionário; e ingestão alimentar, a partir de um questionário de frequência de consumo de alimentos. Os padrões alimentares denominados "dieta mista", "fast food" e "não saudável" foram identificados por meio de análise de componentes principais. Os determinantes de dislipidemia foram examinados através da análise de regressão de Poisson. **Resultados:** A prevalência de dislipidemia, neste estudo, foi de 65,19%. Os pré-escolares que consumiam com menos frequência alimentos do padrão "dieta mista" tiveram um risco maior de concentrações mais altas de lipoproteína de baixa densidade (PR = 2,30; p = 0,004), em comparação com aqueles com um consumo mais frequente do padrão "dieta mista". Os pré-escolares cujas mães apresentaram uma escolaridade mais baixa tiveram um menor risco de concentrações mais altas com lipoproteína de baixa densidade (PR = 0,43; p = 0,003), e os pré-escolares que estavam com sobrepeso/obesos apresentaram maior risco de concentrações mais altas de lipoproteína de baixa densidade (PR = 2,23; p = 0,003).

Conclusão: Os determinantes de dislipidemia identificados neste estudo foram o consumo menos frequente de alimentos do padrão "dieta mista", índice de massa corporal mais alto e um maior nível de escolaridade materna. Este estudo mostra que, apesar da baixa idade do grupo em estudo, já estão apresentando alta prevalência de dislipidemia, que é um fator de risco importante para doença cardiovascular.

© 2013 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

Introduction

Dyslipidemia is a disorder of lipoprotein metabolism that results in elevation of plasma lipid levels, such as high total cholesterol (TC), high low-density lipoprotein cholesterol (LDL-c), low high-density lipoprotein cholesterol (HDL-c), and high triacylglycerol (TAG).¹ In children and adolescents, dyslipidemia is also defined as having a TC, LDL-c, and/or TAG level higher than the 95th percentile or an HDL-c level lower than the 10th percentile for age and gender.²

The prevalence of dyslipidemia in children and adolescents has been high in most countries. According to Al-Shehri,³ dyslipidemia's prevalence varies worldwide from 2.9% to 33% when defining the disease as having a TC level above 200 mg/dL. Studies conducted with children and adolescents in Brazil between 2000 and 2010 observed prevalence rates from 10% to 60.6%.⁴⁻¹⁰

Abnormal levels of lipids and lipoproteins are associated with indirect markers of atherosclerosis, including endothelial dysfunction assessed by flow-mediated dilation in the brachial artery and increased carotid intima-media thickness (cIMT).¹¹ Pediatric elevations of TC, LDL-c, TAG, and the ratio of TC/HDL-c are correlated with increased cIMT and coronary artery calcium during adulthood.¹²

The majority of the cases of dyslipidemia in children and adolescents are related to obesity,¹³⁻¹⁷ high waist-to-hip ratio, family history of dyslipidemia,¹⁷ and lifestyle factors such as high consumption of unhealthy dietary patterns,¹⁸⁻²⁰

meals rich in cholesterol and energy²¹ and low intake of unsaturated fatty acids.²²

Research into determinants of dyslipidemia in Brazilian preschoolers that includes dietary patterns as an explanatory variable has not been conducted yet. Thus, the aim of the present study was to investigate the determinants of dyslipidemia in preschoolers by including dietary patterns, since they can predict disease risk better than single foods or nutrients. The use of dietary patterns makes it possible to identify the cumulative effect of various nutrients on health.

Methods**Type of study and location**

This study used a cross-sectional design nested in a cohort of children who were born in the urban area of the city of Diamantina, Minas Gerais, Brazil, and resided there between September of 2004 and July of 2005.²³ This study²³ aimed to follow the growth and development of this cohort in their first year of life. The newborns' parents were contacted in their homes during the first weeks of life. The recruitment was conducted using the Statement of Live Birth recorded by one of the two hospitals in the city of Diamantina.

Diamantina is a municipality located in the Vale do Jequitinhonha in Minas Gerais, Brazil, that has a mortality rate of 32.8 deaths per 1,000 births among children under

one year old, a literacy rate of 83.4%, a human development index (HDI) of 0.748, and an income HDI of 0.752.²⁴

Subjects and study protocol

At the beginning of the present study, all children were aged 5 years \pm 5 months. They were recruited after an informed consent was obtained from their parents/guardians. This research was conducted from July of 2009 to July of 2010, and data collection was conducted by four nutritionists and one student. Each preschooler was visited in his/her home.

Previous work recommended having five individual per food group listed on the food frequency questionnaire (FFQ), in order to identify dietary patterns.²⁵ In the present study, 24 food groups were identified, thus 120 individuals (5 x 24 groups) were needed for this study. Data from 227 preschoolers were obtained, which was considered a satisfactory sample for the present study.

This study aimed to analyze the determinants of dyslipidemia in addition to identifying dietary patterns, so power for the sample was calculated. Power was calculated post-hoc by using a Poisson regression to estimate the parameter risk difference from being overweight/obese, which was 2.11. The power obtained was 99%, using the statistical software G*Power.²⁶

This study was conducted according to the guidelines presented in the Declaration of Helsinki, and all procedures involving human subjects were approved by the Ethics Committee of the Universidade Federal de Minas Gerais (ref. No. ETIC 545/08). An informed consent was obtained from the preschoolers' parents or guardians.

Anthropometric indicator

Each child's anthropometric status was evaluated by measuring weight and height to obtain the body mass index (BMI). Weight was measured utilizing a portable electronic digital scale with a 150 kg capacity that measured increments of 50 g. Height was measured using a portable stadiometer with a degree of precision of 0.1 cm. The procedures adopted for these measurements followed the protocols recommended by Jelliffe.²⁷

These measurements were taken in a laboratory of the Department of Nursing of the Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM) in the morning with fasting children, and all measurements were performed on a single occasion.

The cut-off point of $\geq +1$ z-score identified preschoolers with elevated body mass index (overweight or obese) for their age using BMI/age.²⁸ To identify z-scores for each child, the World Health Organization (WHO) Anthro and Anthro plus softwares, versions 3.0.1 and 1.0.3, respectively, were used (WHO - Geneva).

Dietary patterns

Dietary patterns (DP) were identified using dietary information collected from a food frequency questionnaire (FFQ) created by Sales et al.²⁹ Before its use, a pilot test was conducted to assess the adequacy of the FFQ for the research.

Any foods not mentioned by the preschoolers' families were excluded, and others were added to the FFQ after the pilot. Details of the food groupings and factor analysis used to generate the dietary patterns are described elsewhere.³⁰

The dietary patterns are presented as a discrete variable. First, the sum of the frequency of consumption of foods contained in each group was calculated. Then, participants were categorized by a dichotomous variable (0 or 1) according to whether they had a value above (1) or below (0) the first quartile frequency of consumption for each food group.

Biochemical indicator

Venous blood samples for quantification of serum TC, LDL-c, HDL-c, and TAG were collected by a trained professional in the morning, between 7:00 and 9:00 a.m., following a fasting period of 10 to 12 hours. Biosafety principles were observed for this collection.

The values for TC, HDL-c, and TAG were obtained utilizing a Cobas Mira Plus device, and were analyzed via absorption photometry using the enzymatic method. LDL-c was determined through calculation using the Friedwald formula while observing the limitations of this methodology.

The cut-offs used to assess the serum levels of TC, TAG, LDL-c, and HDL-c were based on national recommendations.³¹ Elevated levels of TC and LDL-c were defined as ≥ 170 mg/dL and ≥ 130 mg/dL, respectively. TAG levels ≥ 130 mg/dL were considered elevated, and HDL-c levels < 45 mg/dL were considered low. Dyslipidemia was defined as the preschooler presenting abnormal levels of any of these parameters.

Other evaluations

Information regarding factors possibly associated with alterations in the lipid profile was obtained from a questionnaire completed by the child's mother or caregiver at home. This questionnaire provided information about the family's monthly income, maternal education, and behavioral characteristics of the child, such as time spent watching TV.

Statistical analysis

Poisson regression was used to evaluate determinants of dyslipidemia. The variables that presented a p-value < 0.2 in the bivariate analysis were included in a multivariate analysis, and a p-value < 0.05 was considered to be associated with dyslipidemia in the multivariate analysis. This analysis followed a hierarchical approach for the determination of significant factors³² (Fig. 1). The database was constructed in Excel. Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) (SPSS Inc. - Chicago, IL, USA) for Windows, version 19.0.

Results

Of the 227 preschoolers evaluated, 147 (64.7%) presented with dyslipidemia. Among those, 121 (81.7%) had low levels of HDL-c. High levels of TC, LDL-c, and TG were present in




Figure 1 Hierarchical framework of association of dietary patterns with dyslipidemia in preschoolers.

73 (49.3%), 45 (30.4%), and six (2.6%) preschoolers, respectively.

Three dietary patterns were identified: 'mixed diet' consisted of food groups typical of a Brazilian diet; 'snack' consisted of baked food groups that usually do not require preparation for consumption; and 'unhealthy' consisted of sweets and foods rich in lipids and carbohydrates. The 'mixed diet' pattern explained a higher percentage of variance, and was the pattern that best represented the food intake of the sample analyzed.

Table 1 shows the crude prevalence ratios for dyslipidemia according to the socioeconomic, anthropometric, and behavioral characteristics and dietary patterns. This table demonstrates that LDL-c was the lipid associated with the greatest number of variables, followed by HDL-c and TC. Isolated hypertriglyceridemia was not associated with any variable studied, and the results were not presented.

It is worth mentioning that in the 'mixed diet' pattern, which can be considered protective against alteration of the lipid profile, 'low intake' was categorized as a 1. For the other patterns, which can be considered higher risk for alterations of the lipid profile, 'high intake' was categorized as 1.

The regression analysis, adjusted for factors associated with dyslipidemia, is presented in Table 2. Considering the hierarchical framework proposed, which examines the determinants of dyslipidemia among the studied preschoolers, two models were extracted. In model 1, a significant association was found for higher levels of LDL-c, lower maternal education, and low frequency of intake of the 'mixed diet' dietary pattern.

In model 2, high levels of LDL-c were associated with lower maternal education (PR=0.43; $p=0.003$), higher BMI (PR=2.23; $p=0.003$), and low frequency of intake of the 'mixed diet' dietary pattern (PR=2.30; $p=0.004$). In this analysis, high levels of TC and low HDL-c were not associated with any variable.

Of the 281 children in the original cohort, 232 (82.56%) were included in the study. The loss of 17.33% during follow-up was due to families moving away from the area ($n=37$; 75.51%), incorrect addresses ($n=8$, 16.33%); and family refusal to participate ($n=4$; 8.16%). Five parents did not allow their children to provide blood samples, so this study presents results for the remaining 227 preschoolers.

Discussion

The findings of the present study are important warnings that dyslipidemia is part of a worrying reality that needs to be investigated at an early age, such as during preschool. Furthermore, these findings corroborate studies that have shown an increase in the number of cases of this disorder in Brazilian children and adolescents.⁴⁻¹⁰ It is important to mention that the some publications⁴⁻⁶ about alterations in lipid profiles in Brazilian children and adolescents used cutoff points proposed by the NCEP²¹ and recommendations by the III Brazilian guidelines on dyslipidemia,¹ and others⁷⁻¹⁰ used the 1st Brazilian guideline for prevention of atherosclerosis in childhood and adolescence (IDPAIA)³¹

Some international publications on this topic have described how alterations in the lipid profiles of children are also a reality in other countries,^{13-15,17,33-36} but at much lower prevalence than in research reported in Brazil and in the present study.

In this study, the higher prevalence of alterations in lipid profiles was found to be associated with low levels of HDL-c and high levels of LDL-c. These lipoproteins are involved, respectively, in the protection against and formation of atherosclerosis. Individuals with low levels of HDL-c have a higher risk of developing atherosclerosis because this lipoprotein is responsible for various actions that contribute to protection against atherosclerosis.¹ LDL-c is the main atherogenic lipoprotein; its oxidation is the key mechanism in the pathophysiology of an atheroma. Individuals with high levels of LDL-c may present with early clinical manifestations of atherosclerosis.

It is worth mentioning that the lipid profile of children in this study was classified according to the category 'increased' on the IDPAIA,³¹ which favored a higher prevalence of low HDL-c. The IDPAIA uses a raised cutoff point for this lipoprotein. This cutoff point may be inappropriate for this age group, since high levels of HDL-c occur in the presence of increased consumption of fruits and vegetables and physical activity, which are characteristics that are difficult to achieve in this age group.

Among the variables evaluated, three were associated with dyslipidemia (elevated levels of LDL-c) in preschoolers: less frequent consumption of the 'mixed diet' dietary pattern, overweight/obesity, and having a mother with

Table 1 Crude prevalence ratios (PR) and respective p-values of variables associated with altered lipid profiles of preschoolers from Diamantina, Minas Gerais, Brazil.

Variables	PR crude TC	p-value	PR crude LDL-c	p-value	PR crude HDL-c	p-value
<i>Gender</i>						
Male	1		1		1	
Female	1.38	0.09 ^e	1.59	0.08 ^e	0.88	0.37
<i>Body mass index</i>						
Not high	1		1	0.01 ^f	1	0.65
High	1.31	0.22	2.11		0.93	
<i>Time spent watching TV^a</i>						
< 2 hours	1		1		1	
≥ 2 hours	1.19	0.39	1.45	0.19 ^e	0.76	0.02 ^f
<i>Per capita family income^b</i>						
< 1/2 MW	1		1		1	
≥ 1/2 MW	0.89	0.66	0.63	0.15 ^e	1.76	0.03 ^f
<i>Outside-home working mothers</i>						
No	1		1		1	
Yes	0.83	0.32	0.95	0.85	0.74	0.01 ^f
<i>Mother's years of study^c</i>						
> 9 years	1		1		1	
≤ 9 years	0.70	0.06 ^e	0.53	0.02 ^f	1.31	0.04 ^f
<i>Mixed diet patterns^d</i>						
Intake low	1.10	0.66	1.61	0.08 ^{**}	0.88	0.39
Intake high	1		1		1	
<i>Snack patterns</i>						
Intake low	1		1		1	
Intake high	1.38	0.17 ^e	1.15	0.64	1.07	0.62
<i>Unhealthy patterns</i>						
Intake low	1		1		1	
Intake high	0.93	0.74	1.01	0.97	0.84	0.18 ^e

HDL-c, low high-density lipoprotein cholesterol; LDL-c, high low-density lipoprotein cholesterol; MW, Brazilian minimum wage; TC, total cholesterol.

^a Value refers to the median of time spent watching TV.

^b Value refers to the Brazilian minimum wage of US \$288.1 (per month).

^c Value refers to the median years of schooling.

^d Lower and upper first quartile of the distribution of scores for each dietary pattern considered, respectively.

^e significant at the 20% level for the Poisson regression test.

^f significant at the 5% level for the Poisson regression test.

lower levels of education. The 'mixed diet' pattern represents a balanced diet because it consists of foods from all food groups and follows the principles of a healthy diet. Therefore, less frequent consumption may represent a risk factor for elevated levels of LDL-c. Some foods contained in this dietary pattern admittedly provide greater protection against altered lipid profile,³¹ such as leafy vegetables and fruits. Leafy vegetables ($r=0.656$) and fruits ($r=0.618$) showed higher correlations with this AP, which probably explains why less frequent consumption of this dietary pattern was significantly associated with increased levels of LDL-c.

It is interesting to note that some studies^{18,20,37-39} that evaluated the relation between AP and dyslipidemia found that this problem was associated with more frequent consumption of a 'Western' dietary pattern. This pattern is usually composed of foods like red meat, eggs, refined

grains, cafeteria foods, hamburgers, mayonnaise, biscuits, cakes, pies, chocolates, and soft drinks. In this study, while unhealthy patterns were also identified, they not were associated with dyslipidemia.

The association between overweight/obesity and dyslipidemia has been identified in several studies.^{4,9,10,14-16} The Bogalusa Heart Study,¹⁵ conducted in the United States with children and adolescents, found that obese children had 2.4 times and 7.1 times greater chances of having higher levels of total cholesterol and triglycerides, respectively, than children who were not obese. In a study conducted in Brazil, Coronelli et al.¹⁶ observed that obese children had a 2.17 times greater risk of hypercholesterolemia than non-obese children. Alcântara Neto et al.⁹ observed a significant positive association between dyslipidemia and overweight (OR = 3.40) in children and adolescents in the city of Salvador, Bahia.

Table 2 Adjusted prevalence ratios (PR) and respective p-values of variables associated to the alteration in lipid profile of preschoolers. Diamantina, Minas Gerais, Brazil.

Variables studied	PR adjusted TC	p-value	PR adjusted LDL-c	p-value	PR adjusted HDL-c	p-value
MODEL 1: adjusted for distal level + proximal level						
<i>Mother's years of education^a</i>						
≥ 9 years	-		1		-	
< 9 years			0.46	0.01		
<i>Mixed diet patterns^b</i>						
Low intake	-		2.10	0.007	-	
High intake			1			
MODEL 2: adjusted for distal level + intermediate + proximal level						
<i>Body mass index</i>						
Not high	-		1		-	
High			2.23	0.003		
<i>Mother's years of education^a</i>						
≥ 9 years	-		1		-	
< 9 years			0.43	0.003		
<i>Mixed diet patterns^b</i>						
Low intake	-		2.30	0.004	-	
High intake			1			

HDL-c, low high-density lipoprotein cholesterol; LDL-c, high low-density lipoprotein cholesterol; PR, prevalence ratios; TC, total cholesterol; Trace, variable not associated with lipid abnormality in question.

^a Value refers to the median years of schooling.

^b Lower and upper first quartile of the distribution of scores regarding eating pattern.

However, in a study with schoolchildren also residing in the city of Diamantina, MG, Barbosa et al.¹⁰ observed that the correlation between lipid profile (TC, TG, and HDL-c) and anthropometric and body composition parameters was weak. This finding was despite the fact that TC was positively correlated with body fat percentage, whereas HDL-c was negatively correlated with waist-hip ratio in both boys and girls.

According to Asayama et al.,¹⁴ the association between body mass and dyslipidemia has multiple metabolic causes: insulin resistance, hyperinsulinemia, hyperglycemia, and increased protein for transferring cholesterol esters secreted by adipocytes, among other factors.

Another variable associated with dyslipidemia in this study was low maternal education, which exerted a protective effect. This result can be contradictory if it is considered that mothers with lower education may have lower income and less access to information about a balanced diet, which may result in leaving their children more exposed to an unhealthy diet. However, it was observed that the preschoolers with mothers with lower education more often consumed the 'mixed diet' dietary pattern, and this pattern was considered protective against dyslipidemia.

In addition, when a more detailed analysis was performed to understand this relationship, it was observed that the factor that is really influencing this analysis was the lower *per capita* income, and whether the child attended school. That is, low maternal education only remained associated with more frequent consumption of the 'mixed diet' dietary pattern for preschool children attending schools (OR = 4.27) and those with lower *per capita* income (OR = 3.56).³⁰ This

means that the children of mothers with lower education who also had lower incomes and were attending school had less access to processed foods high in fats and sugars and less access to fast food, which put children at the highest risk for dyslipidemia. Moreover, children attending schools eat more balanced meals and likely to be exercising more.

There are some limitations to this study. The most important limitation concerns the assessment of the usual food intake of preschoolers and time spent watching television. Obtaining accurate information on these topics was difficult because most mothers worked outside the home (62.5%), and their children attended school (88.79%). It was difficult for mothers to accurately report what their children eat during the day, and their time spent on each activity. However, it is likely that studies with young children should take these factors into consideration.

The determinants of dyslipidemia identified in this study were less frequent consumption of foods in the 'mixed diet' dietary pattern, higher BMI, and lower levels of maternal education. This study shows that, despite the young age of the group under study, they are already presenting a high prevalence of dyslipidemia, which is an important risk factor for cardiovascular disease. This finding indicates the need to stimulate behavioral change, which can include public policies to fight against inappropriate dietary patterns and prevent diseases that accompany these practices. Children should be the target group of these policies because eating habits are formed in childhood. Inadequate habits that are incorporated in this phase of life and that increase in adolescence may augment the appearance of diseases in adulthood.

Funding

Fundação de Amparo à Pesquisa do Estado de Minas Gerais – FAPEMIG (Process number: APQ-00428-08).

Conflicts of interest

The authors declare no conflicts of interest.

References

- Santos RD. Sociedade Brasileira de Cardiologia III Brazilian guidelines on dyslipidemias and guideline of atherosclerosis prevention from atherosclerosis. Department of Sociedade Brasileira de Cardiologia. *Arq Bras Cardiol.* 2001;77:1–48.
- Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents; National Heart Lung, and Blood Institute. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: summary report. *Pediatrics.* 2011;128:S213–56.
- Al-Shehri SN, Saleh ZA, Salama MM, Hassan YM. Prevalence of hyperlipidemia among Saudi school children in Riyadh. *Ann Saudi Med.* 2004;24:6–8.
- Giuliano I de C, Coutinho MS, Freitas SF, Pires MM, Zunino JN, Ribeiro RQ. Serum lipids in school kids and adolescents from Florianópolis, SC Brazil - Healthy Floripa 2040 study. *Arq Bras Cardiol.* 2005;85:85–91.
- Ribeiro RQ, Lotufo PA, Lamounier JA, Oliveira RG, Soares JF, Botter DA. Additional cardiovascular risk factors associated with excess weight in children and adolescents: the Belo Horizonte heart study. *Arq Bras Cardiol.* 2006;86:408–18.
- Nobre LN, Sarmour SN, Costa Sobrinho PS, Elias FC, Cavaca SC, Trindade R, et al. Lipid profile and weight excess among school children. *Rev Med Minas Gerais.* 2008;18:252–9.
- Pereira PB, Arruda IK, Cavalcanti AM, Diniz A da S. Lipid profile of schoolchildren from Recife, PE. *Arq Bras Cardiol.* 2010;95:606–13.
- Araki MV, Barros C, Santos EG. Análise do perfil lipídico de crianças e adolescentes do estado de Sergipe. *Scientia Plena.* 2010;6:1–6.
- Alcântara Neto OD, Silva RC, Assis AM, Pinto E de J. Factors associated with dyslipidemia in children and adolescents enrolled in public schools of Salvador, Bahia. *Rev Bras Epidemiol.* 2012;15:335–45.
- Barbosa L, Chaves OC, Ribeiro RC. Anthropometric and body composition parameters to predict body fat percentage and lipid profile in schoolchildren. *Rev Paul Pediatr.* 2012;30:520–8.
- Juonala M, Viikari JS, Rönnemaa T, Marniemi J, Jula A, Loo BM, et al. Associations of dyslipidemias from childhood to adulthood with carotid intima-media thickness, elasticity, and brachial flow-mediated dilatation in adulthood: the Cardiovascular Risk in Young Finns Study. *Arterioscler Thromb Vasc Biol.* 2008;28:1012–7.
- Frontini MG, Srinivasan SR, Xu J, Tang R, Bond MG, Berenson GS. Usefulness of childhood non-high density lipoprotein cholesterol levels versus other lipoprotein measures in predicting adult subclinical atherosclerosis: the Bogalusa Heart Study. *Pediatrics.* 2008;121:924–9.
- Cowin I, Emmett P. Cholesterol and triglyceride concentrations, birthweight and central obesity in pre-school children ALSPAC Study Team. *Avon Longitudinal Study of Pregnancy and Childhood. Int J Obes Relat Metab Disord.* 2000;24:330–9.
- Asayama K, Hayashibe H, Dobashi K, Uchida N, Nakane T, Kodera K, et al. Increased serum cholesteryl ester transfer protein in obese children. *Obes Res.* 2002;10:439–46.
- Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics.* 1999;103:1175–82.
- Coronelli CL, de Moura EC. Hypercholesterolemia and its risk factors among schoolchildren. *Rev Saude Publica.* 2003;37:24–31.
- Liao Y, Liu Y, Mi J, Tang C, Du J. Risk factors for dyslipidemia in Chinese children. *Acta Paediatr.* 2008;97:1449–53.
- Dishchekenian VR, Escrivão MA, Palma D, Ancona-Lopez F, Araújo EA, Taddei JA. Dietary patterns of obese adolescents and different metabolic effects. *Rev Nutr.* 2011;24:17–29.
- Northstone K, Emmett P. Multivariate analysis of diet in children at four and seven years of age and associations with socio-demographic characteristics. *Eur J Clin Nutr.* 2005;59:751–60.
- Ambrosini GL, Huang RC, Mori TA, Hands BP, O’Sullivan TA, de Klerk NH, et al. Dietary patterns and markers for the metabolic syndrome in Australian adolescents. *Nutr Metab Cardiovasc Dis.* 2010;20:274–83.
- American Academy of Pediatrics. National Cholesterol Education Program: report of the Expert Panel on Blood Cholesterol Levels in Children and Adolescents. *Pediatrics.* 1992;89:525–84.
- Ribeiro KC, Shintaku RC. The influence of the lipids of the diet on atherosclerosis. *ConScientiae Saúde.* 2004;3:73–83.
- Lessa AC. Alimentação e crescimento no primeiro ano de vida: um estudo de coorte [tese]. Salvador: Universidade Federal da Bahia; 2010.
- Wikipedia. Diamantina. [cited 01 Oct 2012]. Available from: <http://pt.wikipedia.org/wiki/Diamantina>
- Pestano MH, Gageiro JN. Análise de dados para ciências sociais: a complementaridade do SPSS. Lisboa: Edições Sílabo; 2005.
- Portney LG, Watkins MP. Foundations of clinical research: applications to practice. 3rd ed. Upper Saddle River, NJ: Prentice-Hall; 2008.
- Jelliffe DB. Evaluación del estado de nutrición de la comunidad. Geneva: World Health Organization; 1968.
- World Health Organization (WHO). WHO Multicentre Growth Reference Study Group. WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: methods and development. Geneva: World Health Organization. Department of Nutrition for Health and Development. 2006.
- Sales RL, Silva MM, Costa NM, Euclides MP, Eckhardt VF, Rodrigues CM, et al. Development of a questionnaire to assess food intake of population groups. *Rev Nutr.* 2006;19:539–52.
- Nobre LN, Lamounier JA, Franceschini SC. Preschool children dietary patterns and associated factors. *J Pediatr (Rio J).* 2012;88:129–36.
- Back Giuliano I de C, Caramelli B, Pellanda L, Duncan B, Mattos S, Fonseca FH, et al. I guidelines of prevention of atherosclerosis in childhood and adolescence. *Arq Bras Cardiol.* 2005;85:4–36.
- Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol.* 1997;26:224–7.
- Bertrais S, Balkau B, Charles MA, Vol S, Calvet C, Tichet J, et al. Puberty-associated differences in total cholesterol and triglyceride levels according to sex in French children aged 10-13 years. *Ann Epidemiol.* 2000;10:316–23.
- Fesharakinia A, Zarban A, Sharifzadeh GR. Lipid profiles and prevalence of dyslipidemia in schoolchildren in south Khorasan Province, eastern Iran. *Arch Iran Med.* 2008;11:598–601.
- Duarte JA, Ribeiro JC, Oliveira J, Mota J. The relationship between physical activity cholesterol levels in children and adolescents. *Rev Bras Saude Matern Infant.* 2004;4:185–92.
- Ford ES, Li C, Zhao G, Mokdad AH. Concentrations of low-density lipoprotein cholesterol and total cholesterol among

- children and adolescents in the United States. *Circulation*. 2009;119:1108–15.
37. Berg CM, Lappas G, Strandhagen E, Wolk A, Torén K, Rosengren A, et al. Food patterns and cardiovascular disease risk factors: the Swedish INTERGENE research program. *Am J Clin Nutr*. 2008;88:289–97.
 38. Shang X, Li Y, Liu A, Zhang Q, Hu X, Du S, et al. Dietary pattern and its association with the prevalence of obesity and related cardiometabolic risk factors among Chinese children. *PLoS One*. 2012;7:e43183.
 39. Deshmukh-Taskar PR, O'Neil CE, Nicklas TA, Yang SJ, Liu Y, Gustat J, et al. Dietary patterns associated with metabolic syndrome, sociodemographic and lifestyle factors in young adults: the Bogalusa Heart Study. *Public Health Nutr*. 2009;12:2493–503.