



ORIGINAL ARTICLE

Placental and colostral transfer of antibodies reactive with enteropathogenic *Escherichia coli* intimins α , β , or γ ^{☆,☆☆}



Silvia P.N. Altman^{a,b}, Milene Tino-De-Franco^a, Cristiane B. Carbonare^a,
Patricia Palmeira^c, Solange B. Carbonare^{a,*}

^a Instituto Butantan, Laboratório de Imunogenética, São Paulo, SP, Brazil

^b Hospital Israelita Albert Einstein, São Paulo, SP, Brazil

^c Hospital das Clínicas, Instituto da Criança, Laboratório de Investigação Médica (LIM-36), São Paulo, SP, Brazil

Received 21 September 2016; accepted 29 December 2016

Available online 19 March 2017

KEYWORDS

Enteropathogenic
Escherichia coli;
Diarrheagenic
Escherichia coli;
Intimin;
Maternally acquired
immunity;
Immunoglobulin G;
Colostrum

Abstract

Objective: Intimins are protein adhesins of enteropathogenic *Escherichia coli* and enterohemorrhagic *E. coli* capable of inducing attachment and effacement lesions in enterocytes. Anti-intimin antibodies are important for the protection from enteropathogenic *E. coli* and enterohemorrhagic *E. coli* infections because these antibodies inhibit bacterial adhesion and impair the initial step of the pathogenesis. We studied the transfer of maternal anti-intimin antibodies from healthy Brazilian mothers to their newborns through the placenta and colostrum. **Methods:** Serum immunoglobulin G and secretory immunoglobulin A antibodies against conserved and variable regions of intimins α , β , and γ were analyzed using an enzyme linked-immunosorbent assay in the blood and colostrum from 45 healthy women as well as cord blood serum samples from their newborns.

Results: The concentrations of antibodies reactive with α intimin were significantly lower than those of anti- γ and anti-conserved intimin antibodies in the colostrum samples. IgG serum antibodies reactive with all the subtypes of intimins were transferred to the newborns, but the concentrations of anti-conserved intimin serum antibodies were significantly higher in mothers and newborns than concentrations of antibodies against variable regions. The patterns of IgG transfer from mothers to newborns were similar for all anti-intimin antibodies. These values are similar to the percentage transference of total IgG.

[☆] Please cite this article as: Altman SP, Tino-De-Franco M, Carbonare CB, Palmeira P, Carbonare SB. Placental and colostral transfer of antibodies reactive with enteropathogenic *Escherichia coli* intimins α , β , or γ . J Pediatr (Rio J). 2017;93:568–75.

^{☆☆} Study conducted at Instituto Butantan, São Paulo, SP, Brazil.

* Corresponding author.

E-mail: carbosol@usp.br (S.B. Carbonare).

PALAVRAS-CHAVE

Escherichia coli
enteropatogênica;
Escherichia coli
diarreio gênica;
Intimina;
Imunidade materna
adquirida;
Imunoglobulina G;
Colostro

Conclusions: Anti-intimin antibodies are transferred from mothers to newborns through the placenta, and reinforce the protection provided by breastfeeding against diarrheagenic *E. coli* infections.

© 2017 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Transferência placentária e colostrada de anticorpos reativos a *Escherichia coli* enteropatogênica com expressão das intiminas α , β ou γ

Resumo

Objetivo: As Intiminas são adesinas proteicas de *Escherichia coli* enteropatogênicas e enterohemorrágicas capazes de induzir as lesões “attaching and effacing” nos enterócitos. Anticorpos anti-intiminas são importantes para a proteção contra infecções por *E. coli* enteropatogênica e *E. coli* enterohemorrágica porque estes anticorpos inibem a adesão bacteriana impedindo o passo inicial do mecanismo patogênico destas bactérias. Nós estudamos a transferência de anticorpos maternos anti-intiminas de mães brasileiras saudáveis para os seus recém-nascidos através da placenta e do colostro.

Métodos: Anticorpos séricos da classe IgG e secretórios da classe IgA reativos com as porções conservada e variáveis das intiminas α , β e γ foram analisados pelo teste imunoenzimático no sangue e no colostro de 45 parturientes saudáveis e no sangue de cordão umbilical dos seus respectivos recém-nascidos.

Resultados: As concentrações de anticorpos reativos com intimina $\nu\alpha$ foram significativamente mais baixas que as dos anticorpos anti- γ e anti-intimina conservada nas amostras de colostro. Anticorpos IgG séricos reativos com todas as intiminas foram transferidos para os recém-nascidos, mas as concentrações de anti-intimina conservada foram significativamente mais altas tanto nas mães como nos recém-nascidos do que os anticorpos reativos com as regiões variáveis das intiminas. O padrão de transferência de IgG das mães para os recém-nascidos foi muito semelhante para todos os anticorpos anti-intiminas. Os valores de porcentagem de transferência foram semelhantes à transferência de IgG total.

Conclusões: Anticorpos anti-intimina são transferidos das mães para os recém-nascidos pela placenta e corroboram a proteção contra infecções por *Escherichia coli* diarreio gênicas conferida pelo aleitamento materno.

© 2017 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In the developing world, the morbidity and mortality related to pediatric bacterial diarrhea are largely attributed to the diarrheagenic form of *Escherichia coli* (DEC). Newborns and infants up to 1 year of age are particularly vulnerable to diarrhea caused by DEC when they have not been breastfed.¹

Epidemiological studies among Brazilian children have revealed the presence of DEC strains in feces not only of children with diarrhea but also in those of healthy children. Typical (tEPEC) and atypical (aEPEC) forms of enteropathogenic *E. coli* have been detected among these isolates from children in both rural and metropolitan areas.^{2,3} Enterohemorrhagic *E. coli* (EHEC) is an enteric pathogen that is genetically and phenotypically related to EPEC, but distinguishable by its toxin production. Both kinds of bacterial strains produce attaching and effacing lesions (A/E) in the gut mucosa, resulting in intimate contact of the bacteria with the host cell. This phenomenon leads to protein phosphorylation, cell membrane destruction, and expression of bacterial genes, clustered in a genomic island

called locus of enterocyte effacement (LEE).¹ One of the most important proteins encoded by LEE is intimin, a 94-kDa outer-membrane protein involved in intimate bacterial-cell attachment. The intimin molecule is composed of 939 amino acid residues (aa) and has two functional regions: the N-terminal portion, which is highly conserved and is inserted into the bacterial external membrane, and the C-terminal 280 aa portion, which is variable and whose polymorphisms determine the various intimin subtypes.^{4,5} At least 27 subtypes of intimin have been described, but only a few have been implicated in a disease in humans. The prevalence of intimin subtypes varies among tEPEC, aEPEC, and EHEC, in different geographic regions and periods, as demonstrated by epidemiological studies. Intimins α , β , and γ were among the prevalent intimin subtypes in Brazil in the early 2000.^{1,6-8}

Some studies have identified anti-intimin antibodies in the serum of children and adults infected with EPEC or EHEC.^{9,10} The authors have identified anti-intimin antibodies in the serum of healthy adults and children,¹¹⁻¹³ as well as in the colostrum of healthy Brazilian mothers.¹⁴⁻¹⁸ Similar

findings have been described for American and Mexican women.¹⁹ Nonetheless, the transfer of maternal antibody subtypes reactive with intimin to newborns through the placenta has not yet been described.

Antibodies from colostrum inhibit bacterial adhesion to HEp-2 cells *in vitro*; the same anti-EPEC antibodies were found in the feces of breastfeed newborns as present in their mother's colostrum,¹⁵ pointing to a possible mechanism by which breastfeeding confers protection onto newborns against infections caused by EPEC and EHEC.^{14,16,17} In addition, secretory and serum anti-intimin antibodies have been shown to inhibit bacterial adhesion to cultured cells and have been implicated in protection against infection^{15,17,20–22}; therefore, the transfer of anti-intimin antibodies could protect the fetus and newborn against infection. In the present study, the authors assessed the presence of anti-intimin antibodies in the serum and colostrum from healthy mothers and in the umbilical cord of their newborn babies using purified proteins obtained from recombinant bacteria expressing the conserved (cons) and variable regions of intimins α ($v\alpha$), β ($v\beta$), and γ ($v\gamma$).⁴ By simultaneously measuring concentrations of these antibodies in serum samples obtained from the mothers and babies, their possible placental transfer was also assessed.

Materials and methods

Biological samples from human subjects

This study's protocol was approved by the ethics committee of the Hospital Israelita Albert Einstein (HIAE) and of the Universidade de São Paulo, Brazil (CEP number: 06/434); all procedures were in accordance with the 1964 Helsinki declaration, and a written informed consent was obtained from all the participating mothers.

The samples were obtained at HIAE, a private Hospital serving families of a medium to high socioeconomic level, from July to October 2006. Serum, colostrum, and cord blood serum were obtained from 45 healthy mothers and their healthy newborn babies, corresponding to 37 cesarean deliveries. The inclusion criteria were as follows: healthy mothers, without any pathologies recorded during pregnancy or labor, and with negative results on serological tests for HIV, HTLV I/II, hepatitis B and C, Chagas disease, syphilis, toxoplasmosis, and cytomegalovirus. Maternal and cord blood were obtained immediately after delivery and colostrum samples were collected up to 72 h after delivery. All mothers were aged between 22 and 36 years (mean age 31.2), and the total serum immunoglobulin G (IgG) ranged from 508.8 to 1389.1 mg/dL (mean 800.3 mg/dL). The mothers were all well-nourished, predominantly primiparous, with medium to high educational and socioeconomic status, living in good sanitary conditions in the city of São Paulo. The newborn babies were healthy, with adequate weight for the gestational age, 37–41 gestational weeks (mean 38), weighting between 2630 and 3945 g (mean 3298 g).

A control colostrum pool was prepared with equal volumes of 20 samples collected from healthy women, and a control serum pool was prepared with equal volumes of 100 blood samples collected from healthy adult donors. The total secretory immunoglobulin A (SIgA) concentration of

the colostrum pool was determined by a capture enzyme linked-immunosorbent assay (ELISA) using a commercially available standard (Sigma, USA). A serum pool with a known immunoglobulin G (IgG) concentration served as a standard. Samples with low and high concentrations of antibodies were included in each ELISA plate as an internal control of the assay.

Antigens

Purified intimins were obtained from cultures of recombinant bacteria expressing each of the conserved and variable regions of intimins α , β , and γ , as previously described.¹⁸ Briefly, the recombinant *E. coli* M15 bacteria transformed with plasmids pFLV α , pFLV β , pFLV γ , or pFLcons were analyzed by PCR to confirm the presence of the insert after cloning of the plasmid vector. Fragments of the *eae* gene, which encodes intimin, were sequenced for comparison with standard strains registered in GenBank. Protein expression was induced by culturing the bacteria with isopropyl- β -D-thiogalactopyranoside, followed by extraction in a French press and centrifugation. The proteins were finally purified by immobilized metal (Ni) ion adsorption chromatography (Qiagen, USA) and the products were analyzed by sodium dodecyl sulphate polyacrylamide gel electrophoresis.

Anti-intimin antibodies

The concentrations of anti-intimin serum IgG and secretory IgA (SIgA) were determined by ELISA based on the Fomsgaard technique.²³ Concentrations of specific anti-intimin antibodies were determined in comparison with total IgG and IgA concentrations, as described elsewhere.¹⁸ For standardization, half of a microtiter 96-well ELISA plate was coated with anti-IgG as a capture antibody (2.5 μ g/mL) and the other half was coated with each of the purified intimins obtained from recombinant bacteria expressing cons, $v\alpha$, $v\beta$, and $v\gamma$ ⁴ at the same concentration (2.5 μ g/mL). Appropriate serial dilutions of the control serum pool (with known total IgG concentration determined by nephelometry) were added on both sides of the plate. The assay was developed with an anti-IgG conjugate. The optical density values were established and the concentration of anti-intimin antibodies was determined by comparison with the standard curve of total IgG. This standard procedure was also performed for SIgA in a colostrum pool. Then, the individual serum and colostrum samples in serial dilutions starting from 1:10 were analyzed in ELISA plates coated with the recombinant intimins, with the serum or colostrum pool used to construct the standard curve. Samples with high and low concentrations of antibodies were included in each plate as an internal control. The total IgG concentrations in maternal and cord serum samples were measured by an immunoturbidimetric assay (Abbott Diagnostics, USA). The placental transfer percentages of the total and intimin-specific IgG antibodies were defined in each assay as follows:

$$\%T = \left(\frac{\text{concentration of antibodies in cord serum}}{\text{concentration of antibodies in the maternal serum}} \right) \times 100$$

Statistical analysis

All results were subjected to statistical analysis including descriptive analyses with box plot graphs, non-parametric ANOVA with Friedman's multiple comparison test, and the Spearman correlation test, at a 95% confidence level, using the GraphPad Prism software (Graph Pad Software Inc., USA).

Results

Fig. 1 shows the concentrations of antibodies reactive with the four purified recombinant antigens (cons, $v\alpha$, $v\beta$, and $v\gamma$) present in 45 colostrum samples, 45 maternal serum samples, and 45 cord serum samples from their respective newborns. The box plots show the maximal and minimal values, 25% and 75% percentiles, medians, and means.

The analyses of colostrum samples by the Friedman test for multiple comparisons showed that the concentrations of anti- $v\alpha$ IgA antibodies were significantly lower than those of anti- $v\gamma$ and anti-cons, but there were no statistically significant differences within the other pairs of values (Fig. 1A). The concentrations of IgA antibodies varied widely, and the means above the medians indicated that it was not a normal distribution.

The comparison of specific IgG revealed that the concentration of anti-cons was significantly higher when compared with the other subtypes in the mothers' and newborns' serum samples. The anti- $v\alpha$ antibody levels were higher than those of anti- $v\beta$, but there were no significant differences between these two subtypes and anti- $v\gamma$ antibodies (Fig. 1B and C). These results were confirmed by the frequency distribution analysis (data not shown).

Fig. 2 shows that the IgG transfer from mothers to newborns was very similar for all the antibody subtypes analyzed. The mean transference percentage of anti-intimin antibodies ranged from 142% to 149% and the medians, from 130% to 146%. These values are similar to the percentage of transference of total IgG (mean = 143%, median = 137%). There were no significant differences among the transference percentages of antibodies reactive with the various intimin subtypes.

Table 1 shows results of the Spearman correlation tests. In maternal serum samples, there was a significant correlation between anti- $v\alpha$ and anti- $v\beta$ antibodies ($p < 0.0001$). There was no statistically significant correlation between $v\gamma$ and cons. The same pattern of correlation was observed in newborns' serum samples. The correlation coefficients were high in all pairs analyzed for the colostrum samples.

Table 2 shows significant correlation coefficients for all intimin types between maternal and newborn serum samples; therefore, the transference of antibodies from mothers to newborns was equally efficient for all the subtypes of intimin. Low correlation coefficients between maternal serum IgG and colostrum SIgA antibodies were observed, as the generation of serum IgG and secretory IgA antibodies proceeds *via* independent mechanisms.

Discussion

In this study, the authors provided evidence of placental and colostrum transfer of anti-EPEC and anti-EHEC antibodies to




Figure 1 Box-plot distributions of samples by their anti-intimin concentrations: maximal and minimal values (top and bottom lines outside the box, respectively), means (\bullet) and medians (\blacktriangle), and 75% and 25% quartiles (top and bottom sides of the box, respectively). (a) SIgA antibodies in colostrum samples. (b) IgG antibodies in maternal serum samples. (c) IgG antibodies in newborns' serum samples. Significant differences: * $p < 0.05$, ** $p < 0.01$.

Table 1 Correlation coefficients of the concentrations of anti-intimin antibodies in serum (IgG) and colostrum (SIgA) samples and IgG transference percentage from mothers to newborns, in pairs of intimin types.

Anti-intimin antibodies	Maternal sera	Newborn sera	Colostrum	IgG transference (%)
	IgG	IgG	SIgA	
v α vs. v β	0.8014 <i>(0.0001)</i>	0.8132 <i>(0.0001)</i>	0.8235 <i>(0.0001)</i>	0.3447 <i>(0.0204)</i>
v α vs. v γ	0.3590 <i>(0.0154)</i>	0.4586 <i>(0.0015)</i>	0.7768 <i>(0.0001)</i>	0.4603 <i>(0.0015)</i>
v α vs. cons	0.3159 <i>(0.0345)</i>	0.5995 <i>(0.0001)</i>	0.7809 <i>(0.0001)</i>	0.3121 <i>(0.0369)</i>
v β vs. v γ	0.4433 <i>(0.0023)</i>	0.5034 <i>(0.0004)</i>	0.7232 <i>(0.0001)</i>	0.5306 <i>(0.0002)</i>
v β vs. cons	0.3495 <i>(0.0186)</i>	0.4934 <i>(0.0006)</i>	0.7111 <i>(0.0001)</i>	0.2462 <i>(0.1031)</i>
v γ vs. cons	0.1043 <i>(0.4954)</i>	0.2221 <i>(0.1426)</i>	0.7628 <i>(0.0001)</i>	0.3306 <i>(0.0265)</i>

v α , variable region of intimin α ; v β , variable region of intimin β ; v γ , variable region of intimin γ ; cons, conserved region of intimin. The coefficients were determined by the Spearman correlation test, at a 95% confidence level. *p*-Values are shown in italics; *p*-values < 0.0001 are highlighted in bold.

newborn babies; these antibodies are likely to confer protection onto the neonates against diarrhea caused by these pathogens. Anti-intimin antibodies can inhibit the bacterial adhesion to epithelial cells, and this ability has been demonstrated for SIgA antibodies present in human colostrum and milk and serum antibodies from humans and immunized animals.^{13–17,22,24} Based on this property, intimins have been proposed as a potential candidate for a vaccine against EPEC and EHEC infection.^{20,21}

Antibodies against the conserved and variable regions of intimins α , β , and γ were found in all serum and colostrum samples analyzed, probably due to the wide prevalence of EPEC in the Brazilian population. This study was conducted




Figure 2 Transfer of IgG anti-intimin antibodies and total IgG from mothers to newborns. Box-plot distribution of samples by their percentages of transference of anti-v α , anti-v β , anti-v γ , and anti-cons and transfer of total IgG: maximal and minimal values (top and bottom lines outside the box, respectively), means (●) and medians (▲), and 75% and 25% quartiles (top and bottom sides of the box, respectively).

Table 2 Correlation coefficients of anti-intimin antibody concentrations between maternal serum (IgG) and colostrum (SIgA) samples, and between maternal and cord serum samples (IgG).

Intimin	Maternal serum vs. colostrum	Maternal serum vs. newborn serum
v α	0.0028 <i>(0.9853)</i>	0.8727 (0.0001)
v β	0.1475 <i>(0.3337)</i>	0.9568 (0.0001)
v γ	0.1743 <i>(0.2578)</i>	0.9320 (0.0001)
cons	0.2265 <i>(0.1346)</i>	0.8519 (0.0001)

v α , variable region of intimin α ; v β , variable region of intimin β ; v γ , variable region of intimin γ ; cons, conserved region of intimin.

Coefficients were determined by the Spearman correlation test, at a 95% confidence level. *p*-Values are shown in italics; *p*-values < 0.0001 are highlighted in bold.

using samples collected from mothers from medium to high socioeconomic levels. The authors have demonstrated similar findings when analyzing samples collected from mothers of lower socioeconomic layers and living in poorly sanitized areas, suggesting that EPEC prevalence is independent of the socioeconomic status or hygienic conditions of the patients.^{11,12,16,18} Similar findings were reported by Araujo et al. regarding the detection of DEC in fecal samples of children from poor urban areas and in those of children receiving in high-socioeconomic-level private medical services.²⁵

Healthy Brazilian children develop antibodies reactive to EPEC virulence-associated factors such as intimins.¹¹ By the end of the first year of life, these infants exhibit serum and salivary anti-EPEC antibody repertoires equivalent to those of healthy adults in the same population. Intimins α , β , and γ are shared by EPEC (whether tEPEC or aEPEC) and EHEC strains, and are present in DEC found in domestic and farm animals that are in close contact with people, suggesting

that intimin is a common antigen ubiquitously present in the Brazilian population.²⁶

The anti-intimin antibody subtypes found in the samples analyzed in this study may inhibit bacterial adhesion, because this portion of the intimin molecule is responsible for interactions with the host cells, pointing to a role in protection against the bacteria.²²

The statistical analysis of IgG detected in both maternal and newborn serum samples revealed that concentrations of antibodies against the conserved region of intimin were higher in comparison with antibodies to the variable regions of the subtypes α , β , and γ . The conserved region is common to all subtypes; therefore, it is reasonable to expect that all individuals will develop these antibodies regardless of the specific intimin subtype present in the contacting bacteria. Similar results were observed by Zapata-Quintanilla et al.¹⁸ in their analyses of serum samples from healthy adults (blood donors). Nevertheless, this profile was different in colostrum samples, where the concentration of antibodies reactive to $\nu\beta$, $\nu\gamma$, and cons were all similar, but the anti- $\nu\alpha$ antibody concentration was lower than that of anti- $\nu\gamma$ and anti-cons. This pattern was not identified by Zapata-Quintanilla et al.,¹⁸ who found lower anti- $\nu\gamma$ levels. This discrepancy may be attributable to variations in the antibody repertoires among the mothers, as the characteristics of the donors in the study by Zapata-Quintanilla were different from those of the present study regarding the socio-economic level (public hospital versus private hospital, respectively), age (mean 24.6 vs. 31.2 years), type of delivery (24.4% vs. 82% cesarean section), and period of sample collection (September 1997 to February 1998 vs. July to October 2006). Epidemiological studies have indicated that the prevalence of intimin γ has increased with the emergence of aEPEC in recent years.^{1,3,6-8,25}

Intimins are frequently found in bacteria isolated not only from patients with diarrhea, but also from healthy people and domestic animals in Brazil.^{1,6,7,26} Intimin α was detected in tEPEC and aEPEC strains. Intimin β is associated with tEPEC and aEPEC, as well as with EHEC, whereas intimin γ is found in aEPEC and EHEC. Isolation of these bacterial strains in Brazil can explain the presence of these antibodies in serum and colostrum samples. In spite of the low incidence of EHEC infection among Brazilian adults, anti-intimin γ antibodies are present and show a positive correlation with antibodies reactive with other intimin subtypes. The increasing frequencies of aEPEC strains in the environment may be responsible for the presence of anti-intimin γ antibodies among healthy Brazilians. Recent studies have implicated intimin subtypes β and γ in the induction of diarrhea in humans, and subtype β was found to be significantly more frequent in disease cases when compared with healthy carriers.⁸

There is a significant correlation between the concentration of anti-intimin α and β IgG antibodies in the maternal serum and newborns' serum, which may be due to cross-reactions among IgG antibodies targeting these intimins. Indeed, the gene sequences of the variable portions of these intimins are 65% homologous,¹⁸ which explains the presence of cross-reacting antibodies. Therefore, this phenomenon implies that whatever the intimin subtype candidate for a vaccine, this subtype may raise specific antibodies but can

also induce cross-reactive antibodies directed to the other subtypes.

The concentrations of SIgA antibodies reactive with all types of intimins present in colostrum samples showed significant correlation coefficients, suggesting that there is a certain degree of cross-reaction among these antibodies. This situation may be explained by the elevated homology between the intimin molecules and by the lower specificity of colostrum SIgA antibodies as compared to serum IgG. Nevertheless, the higher concentrations of anti-intimin antibodies in colostrum samples when compared with serum samples support the crucial role of breastfeeding in protection of infants from DEC infection. In humans, milk SIgA antibodies are not significantly absorbed by the mucosa of the breastfed baby, but act as local defense. In a previous study, the authors have shown, using immunoblotting, that the reactivity of anti-EPEC SIgA antibodies of the colostrum samples was the same as the reactivity of anti-EPEC antibodies in the feces of the respective babies. In addition, SIgA antibodies isolated from colostrum by affinity chromatography inhibit EPEC adhesion to HEp-2 cells.¹⁵

The present results indicate that different anti-intimin antibodies are transferred from mothers to newborns with equal efficiency through the placenta. These results support the hypothesis that, in this case, the maternal-fetal transfer is not influenced by antibody specificity. The transference percentages of anti-intimin antibodies in the present study were found to be equivalent to the transference percentage of total IgG, which is approximately 140%. Similar findings have been reported by other authors.^{27,28}

Although there is a high frequency of DEC isolates in Brazil, strains of EHEC are not as common as in other countries. It can be hypothesized that, besides environmental features such as climate and alimentary habits, people may develop EHEC immunity resulting from contact with other strains (e.g., DEC). Some studies revealed the presence of antibodies reactive with various lipopolysaccharide (LPS) serotypes, such as O111, O55, O157, and O26, in serum and colostrum samples from healthy Brazilians.^{12,13} Other studies have shown the maternal-fetal transfer of IgG antibodies reactive with O111, O16, and O6 LPS, as well as O157, to be associated with neonatal sepsis.^{29,30} These LPS antigens are commonly found in different pathotypes of *E. coli* that are agents of intestinal or systemic diseases. Taken together, these observations and the present work allow us to conclude that protection from EHEC can be conferred by contact with EPEC or other *E. coli* strains.

The present results confirm that maternal antibodies reactive with intimins are transferred to the newborns through the placenta, and that the presence of high concentrations of these antibodies in colostrum supports the protective role of breastfeeding.

Funding

The financial support was provided by FAPESP – 03/13250-3.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors would like to thank all the mothers for their consent to donate the samples. They would also like thank the Maternity, the Milk Bank, and the Cryopreservation Laboratory of Hospital Israelita Albert Einstein, for their help with the sample collection; the Brazilian Division of Abbot Diagnosis, for the donation of the Immunoturbidimetric Kit for the analysis of IgG; Dr. Lucy B. Zapata-Quintanilla, for the help with experiments, Rosana Prisco, for the statistical analysis; and Dr. Yanira Rizzo Vasquez, for review of the manuscript and her valuable suggestions (Institute of Pharmaceutical Science – King's College, London, United Kingdom).

References

- Gomes TA, Elias WP, Scaletsky IC, Guth BE, Rodrigues JF, Piazza RM, et al. Diarrheagenic *Escherichia coli*. *Braz J Microbiol*. 2016;47:3–30.
- Lozer DM, Souza TB, Monfardini MV, Vicentini F, Kitagawa SS, Scaletsky IC, et al. Genotypic and phenotypic analysis of diarrheagenic *Escherichia coli* strains isolated from Brazilian children living in low socioeconomic level communities. *BMC Infect Dis*. 2013;13:418.
- Moreno AC, Filho AF, Gomes Tdo A, Ramos ST, Montemor LP, Tavares VC, et al. Etiology of childhood diarrhea in the north-east of Brazil: significant emergent diarrheal pathogens. *Diagn Microbiol Infect Dis*. 2010;66:50–7.
- Adu-Bobie J, Frankel G, Bain C, Goncalves AG, Trabulsi LR, Douce G, et al. Detection of intimin alpha, beta, gamma, and delta, four intimin derivatives expressed by attaching and effacing microbial pathogens. *J Clin Microbiol*. 1998;36:662–8.
- Adu-Bobie J, Trabulsi LR, Carneiro-Sampaio MM, Dougan G, Frankel G. Identification of immunodominant regions within the C-terminal cell binding domain of intimin alpha and intimin beta from enteropathogenic *Escherichia coli*. *Infect Immun*. 1998;66:5643–9.
- Franco RT, Araújo LD, Penna FJ, Magalhães PP, Mendes EN. Intimin subtyping of atypical enteropathogenic *Escherichia coli* isolated from children with and without diarrhea: a possible temporal shift in the distribution of intimin alleles. *Diagn Microbiol Infect Dis*. 2012;74:81–3.
- Pitondo-Silva A, Nakazato G, Falcão JP, Irino K, Martinez R, Darini AL, et al. Phenotypic and genetic features of enteropathogenic *Escherichia coli* isolates from diarrheal children in the Ribeirão Preto metropolitan area, São Paulo State, Brazil. *APMIS*. 2015;123:128–35.
- Hu J, Torres AG. Enteropathogenic *Escherichia coli*: foe or innocent bystander? *Clin Microbiol Infect*. 2015;21:729–34.
- Martinez MB, Taddei CR, Ruiz-Tagle A, Trabulsi LR, Girón JA. Antibody response of children with enteropathogenic *Escherichia coli* infection to the bundle-forming pilus and locus of enterocyte effacement-encoded virulence determinants. *J Infect Dis*. 1999;179:269–74.
- Voss E, Paton AW, Manning PA, Paton JC. Molecular analysis of Shiga toxigenic *Escherichia coli* O111:H⁻ proteins which react with sera from patients with hemolytic-uremic syndrome. *Infect Immun*. 1998;66:1467–72.
- Carbonare CB, Carbonare SB, Carneiro-Sampaio MM. Early acquisition of serum and saliva antibodies reactive to enteropathogenic *Escherichia coli* virulence-associated proteins by infants living in an endemic area. *Pediatr Allergy Immunol*. 2003;14:222–8.
- Palmeira P, Carbonare SB, Guth BE, Carbonare CB, Pontes GN, Tino-De-Franco M, et al. Acquisition of serum antibodies reactive with enterohemorrhagic *Escherichia coli* virulence-associated factors by healthy Brazilian children and adults. *Pediatr Infect Dis J*. 2009;28:1089–94.
- Zapata-Quintanilla LB, Palmeira P, Tino-De-Franco M, Amaral JA, Carbonare CB, Carbonare SB. Systemic antibody response to diarrheagenic *Escherichia coli* and LPS O111, O157 and O55 in healthy Brazilian adults. *Scand J Immunol*. 2006;64:661–7.
- Camara LM, Carbonare SB, Silva ML, Carneiro-Sampaio MM. Inhibition of enteropathogenic *Escherichia coli* (EPEC) adhesion to HeLa cells by human colostrum: detection of specific sIgA related to EPEC outer-membrane proteins. *Int Arch Allergy Immunol*. 1994;103:307–10.
- Carbonare SB, Silva ML, Palmeira P, Carneiro-Sampaio MM. Human colostrum IgA antibodies reacting to enteropathogenic *Escherichia coli* antigens and their persistence in the faeces of a breastfed infant. *J Diarrhoeal Dis Res*. 1997;15:53–8.
- Delneri MT, Carbonare SB, Silva ML, Palmeira P, Carneiro-Sampaio MM. Inhibition of enteropathogenic *Escherichia coli* adhesion to HEp-2 cells by colostrum and milk from mothers delivering low-birth-weight neonates. *Eur J Pediatr*. 1997;156:493–8.
- Palmeira P, Carbonare SB, Amaral JA, Tino-De-Franco M, Carneiro-Sampaio MM. Colostrum from healthy Brazilian women inhibits adhesion and contains IgA antibodies reactive with Shiga toxin-producing *Escherichia coli*. *Eur J Pediatr*. 2005;164:37–43.
- Zapata-Quintanilla LB, Tino-De-Franco M, Nuñez SP, Carbonare CB, Vulcano AB, Carbonare SB. Seric and secretory antibodies reactive to alpha, beta and gamma intimins of *Escherichia coli* in healthy Brazilian adults. *FEMS Immunol Med Microbiol*. 2009;57:32–9.
- Noguera-Obenza M, Ochoa TJ, Gomez HF, Guerrero ML, Herrera-Insua I, Morrow AL, et al. Human milk secretory antibodies against attaching and effacing *Escherichia coli* antigens. *Emerg Infect Dis*. 2003;9:545–51.
- Dean-Nystrom EA, Gansheroff LJ, Mills M, Moon HW, O'Brien AD. Vaccination of pregnant dams with intimin (O157) protects suckling piglets from *Escherichia coli* O157:H7 infection. *Infect Immun*. 2002;70:2414–8.
- Ferreira PC, Campos IB, Abe CM, Trabulsi LR, Elias WP, Ho PL, et al. Immunization of mice with *Lactobacillus casei* expressing intimin fragments produces antibodies able to inhibit the adhesion of enteropathogenic *Escherichia coli* to cultivated epithelial cells. *FEMS Immunol Med Microbiol*. 2008;54:245–54.
- Gansheroff LJ, Wachtel MR, O'Brien AD. Decreased adherence of enterohemorrhagic *Escherichia coli* to HEp-2 cells in the presence of antibodies that recognize the C-terminal region of intimin. *Infect Immun*. 1999;67:6409–17.
- Fomsgaard A. Antibodies to lipopolysaccharides: some diagnostic and protective aspects. *APMIS Suppl*. 1990;18:1–38.
- Amaral JA, Tino De Franco M, Carneiro-Sampaio MM, Carbonare SB. Anti-enteropathogenic *Escherichia coli* immunoglobulin Y isolated from eggs laid by immunised Leghorn chickens. *Res Vet Sci*. 2002;72:229–34.
- Araujo JM, Tabarelli GF, Aranda KR, Fabbricotti SH, Fagundes-Neto U, Mendes CM, et al. Typical enteroaggregative and atypical enteropathogenic types of *Escherichia coli* are the most prevalent diarrhea-associated pathotypes among Brazilian children. *J Clin Microbiol*. 2007;45:3396–9.
- Aidar-Ugrinovich L, Blanco J, Blanco M, Blanco JE, Leomil L, Dahbi G, et al. Serotypes, virulence genes, and intimin types of Shiga toxin-producing *Escherichia coli* (STEC) and enteropathogenic *E. coli* (EPEC) isolated from calves in São Paulo, Brazil. *Int J Food Microbiol*. 2007;115:297–306.

27. Palmeira P, Quinello C, Silveira-Lessa AL, Zago CA, Carneiro-Sampaio M. IgG placental transfer in healthy and pathological pregnancies. *Clin Dev Immunol*. 2012;2012:985646.
28. de Voer RM, van der Klis FR, Nooitgedagt JE, Versteegh FG, van Huiseling JC, van Rooijen DM, et al. Seroprevalence and placental transportation of maternal antibodies specific for *Neisseria meningitidis* serogroup C, *Haemophilus influenzae* type B, diphtheria, tetanus, and pertussis. *Clin Infect Dis*. 2009;49:58–64.
29. Nagao AT, Martinez CC, Vieira VS, Takano OA, Costa-Carvalho BT, Carneiro-Sampaio MM. Placental transfer of IgG and IgG subclass antibodies anti-purified *Escherichia coli* LPS O16, O6 and O111. *Scand J Immunol*. 1998;47:609–14.
30. Palmeira P, Yu Ito L, Arslanian C, Carneiro-Sampaio MM. Passive immunity acquisition of maternal anti-enterohemorrhagic *Escherichia coli* (EHEC) O157:H7 IgG antibodies by the newborn. *Eur J Pediatr*. 2007;166:413–9.