



## ORIGINAL ARTICLE

# Skinfold reference curves and their use in predicting metabolic syndrome risk in children<sup>☆,☆☆</sup>



Alynne C.R. Andaki<sup>a,\*</sup>, Teresa M.B. de Quadros<sup>b</sup>, Alex P. Gordia<sup>b</sup>, Jorge Mota<sup>c</sup>, Adelson L.A. Tinôco<sup>d</sup>, Edmar L. Mendes<sup>a</sup>

<sup>a</sup> Universidade Federal do Triângulo Mineiro (UFTM), Departamento de Ciências do Esporte, Uberaba, MG, Brazil

<sup>b</sup> Universidade Federal do Recôncavo da Bahia (UFRB), Centro de Formação de Professores, Cruz das Almas, BA, Brazil

<sup>c</sup> Universidade do Porto, Faculdade do Desporto, Porto, Portugal

<sup>d</sup> Universidade Federal de Viçosa (UFV), Departamento de Nutrição e Saúde, Viçosa, MG, Brazil

Received 28 July 2016; accepted 16 November 2016

Available online 23 May 2017

## KEYWORDS

Anthropometry;  
Metabolic syndrome  
X;  
Cross-sectional  
studies;  
Child

## Abstract

**Objectives:** To draw skinfold (SF) reference curves (subscapular, suprailiac, biceps, triceps) and to determine SF cutoff points for predicting the risk of metabolic syndrome (MetS) in children aged 6–10 years old.

**Methods:** This was a cross-sectional study with a random sample of 1480 children aged 6–10 years old, 52.2% females, from public and private schools located in the urban and rural areas of the municipality of Uberaba (MG). Anthropometry, blood pressure, and fasting blood samples were taken at school, following specific protocols. The LMS method was used to draw the reference curves and ROC curve analysis to determine the accuracy and cutoff points for the evaluated skinfolds.

**Results:** The four SF evaluated (subscapular, suprailiac, biceps, and triceps) and their sum ( $\sum 4\text{SF}$ ) were accurate in predicting MetS for both girls and boys. Additionally, cutoffs have been proposed and percentile curves (p5, p10, p25, p50, p75, p90, and p95) were outlined for the four SF and  $\sum 4\text{SF}$ , for both genders.

**Conclusion:** SF measurements were accurate in predicting metabolic syndrome in children aged 6–10 years old. Age- and gender-specific smoothed percentiles curves of SF provide a reference for the detection of risk for MetS in children.

© 2017 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<sup>☆</sup> Please cite this article as: Andaki AC, Quadros TM, Gordia AP, Mota J, Tinôco AL, Mendes EL. Skinfold reference curves and their use in predicting metabolic syndrome risk in children. J Pediatr (Rio J). 2017;93:490–6.

<sup>☆☆</sup> Study carried out at Universidade Federal do Triângulo Mineiro, Uberaba, MG, Brazil.

\* Corresponding author.

E-mail: [alynneandaki@yahoo.com.br](mailto:alynneandaki@yahoo.com.br) (A.C. Andaki).

**PALAVRAS-CHAVE**

Antropometria;  
Síndrome X;  
Estudos de corte  
transversal;  
Crianças

**Curvas de referência de dobras cutâneas e sua utilização na predição do risco de síndrome metabólica em crianças****Resumo**

**Objetivos:** Desenhar curvas de referência de quatro dobras cutâneas (subescapular, suprailíaca, bíceps, tríceps) e determinar pontos de corte para predizer o risco de SM em crianças de seis a 10 anos de idade.

**Métodos:** Estudo epidemiológico de base populacional, corte transversal, com amostra probabilística, estratificada por segmento de ensino, com 1.480 crianças de 6 a 10 anos de idade, 52,2% do sexo feminino, oriundas de escolas públicas e privadas, localizadas na zona urbana e rural do município de Uberaba (MG). Antropometria (dobras cutâneas), pressão arterial e as coletas de sangue em jejum foram realizadas em espaço reservado na escola, segundo protocolos específicos. O método LMS foi utilizado para desenhar as curvas de referência e análise de curva ROC para determinar a acurácia e pontos de corte para as dobras cutâneas avaliadas.

**Resultados:** As quatro DC avaliadas (subescapular, suprailíaca, bíceps e tríceps) e o seu somatório ( $\sum 4DC$ ) foram acurados na predição da SM para meninas e meninos. Adicionalmente, pontos de corte foram propostos e curvas percentílicas (p5, p10, p25, p50, p75, p90 e p95) foram delineadas para as quatro DC e o  $\sum 4DC$ , para ambos os sexos.

**Conclusão:** Medidas de DC foram acuradas em predizer SM em escolares de seis a 10 anos de idade. As curvas percentílicas de DC desenhadas por idade e sexo fornecem referência na detecção do risco de SM em crianças.

© 2017 Publicado por Elsevier Editora Ltda. em nome de Sociedade Brasileira de Pediatria. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Cardiovascular diseases are the main cause of death in Brazil<sup>1</sup> and worldwide.<sup>2</sup> Cardiovascular risk factors, such as total and visceral obesity, dyslipidemias, arterial hypertension, hyperglycemia, and hyperinsulinemia have been considered determinants for the development of cardiovascular diseases. Metabolic syndrome (MetS), characterized by the presence of three or more of these risk factors,<sup>3</sup> has gained importance due to its consistent association with cardiovascular morbimortality.<sup>4</sup>

The presence of cardiovascular risk factors and MetS has been observed in adults<sup>5</sup> as well as in children and adolescents.<sup>6</sup> In recent years, concerns about the diagnosis of and early intervention on these metabolic disorders has increased due to evidence that risk factors observed in childhood and adolescence tend to persist and worsen in adulthood.<sup>7</sup>

In addition to blood pressure (BP) and waist circumference measurements, biochemical analyses of blood fractions of HDL-c, triglycerides, and glucose should be performed for the diagnosis of MetS. Blood collection is an invasive, expensive and difficult to perform technique, especially in children and adolescents. In turn, the use of practical and low-cost methods may be an important alternative for the screening of MetS at the population level. In this sense, skinfolds (SFs) were shown to be a promising tool for MetS screening in the pediatric population, due to its strong correlation with subcutaneous adiposity.<sup>8</sup>

According to Ali et al.,<sup>9</sup> the accumulation of subcutaneous adiposity is a strong predictor of insulin resistance and hypertriglyceridemia in children and adolescents, and was a stronger predictor of cardiometabolic risk factors than

visceral fat. Furthermore, the influence of subcutaneous adiposity on MetS risk is present in children and adolescents, but not in adults.<sup>9</sup>

Percentile curves for SFs have been developed with samples of young North-American,<sup>10</sup> Polish,<sup>11</sup> and Indian populations.<sup>12</sup> It is noteworthy that the World Health Organization<sup>13</sup> presented an important publication in 2007, in which reference curves for anthropometric measurements, including triceps and subscapular SFs were created as an international reference for multi-country populations (Brazil, United States, Ghana, India, Norway, and Oman). However, to the best of the authors' knowledge, SF percentile curves have not been proposed for Brazilian children, nor are there any SF percentile curves that can be used to predict MetS in the pediatric population. Reference curves from other populations may not be applicable to Brazilian children due to ethnic, cultural, and socioeconomic differences.

Therefore, the present study aimed to design reference curves of four SFs (subscapular, suprailiac, biceps, and triceps) and to determine cutoffs to predict the risk of MetS in children aged 6–10 years.

## Methods

A cross-sectional, population-based epidemiological study was carried out with a probabilistic sample of children aged 6–10 years from public and private schools located in the urban and rural areas of the city of Uberaba, MG, Brazil.

The sample calculation considered the number of children enrolled in Elementary School (1st to 9th year of schooling), a prevalence of MetS of 50% (unknown prevalence in the municipality), tolerable error of 3.5% and

confidence level of 95%. The minimum sample size was 768 children; after adding 10% to compensate for losses and refusals and 20% to minimize confounding factors, the sample totaled 1014 children.

For sample selection, the schools were stratified according to the type of school as municipal, state, or private.

The World Health Organization<sup>14</sup> recommends that 10–15 sample collection points (schools) be used for epidemiological surveys, and that the number of subjects in each age group should vary between 25 and 50 for each site. Therefore, 15 of the 90 eligible schools of the municipality of Uberaba were randomly selected using the random number table. For the adequacy and representativeness of the local population, the number of children in each stratum was determined proportionally to the number of enrollments, according to data provided by the State Education Secretariat. Municipal schools accounted for 43.6% of enrollments, 41.9% of students were enrolled from state schools and 14.5% from private schools.

After approval by the Ethics Committee in Research with Human Beings of Universidade Federal do Triângulo Mineiro (Protocol CEP/UFTM: 1710), the school principals were contacted to obtain authorization and to schedule the collections. Students who met the inclusion criteria and were interested in participating in the research received the informed consent form for the information and signature of their parents and/or guardians. Anthropometry and blood collections were performed at the school itself, following specific protocols.

The biceps (BSF), triceps (TSF), subscapular (SSSF), and suprailiac (SISF) SFs were obtained using an adipometer (Lange Skinfold Caliper, England, UK) exerting constant pressure of 10 g/mm<sup>2</sup>. Measurements were performed on the right side of the body, with three non-consecutive repetitions for each measurement. The final measure constituted the mean of the three values. All measurements were performed by two qualified evaluators submitted to previous training and calibration. The results were interpreted alone, as well as by the sum of the four SFs assessed, and were expressed in millimeters (mm).

BP was measured using a mercury column sphygmomanometer (Unitec, São Paulo, Brazil), with cuffs of appropriate sizes for the circumference of the children's arms, in accordance with the VI Brazilian Guidelines for Hypertension.<sup>15</sup> Three BP measurements were performed at the first visit, after which the first was discarded and the mean of the last two were considered. When the child had an alteration in systolic or diastolic BP above the 90th percentile, two further measurements were performed on two different days following the same procedure adopted on the first day.<sup>15</sup>

The volunteers were invited to go to school, following a 12-h fast, on certain days and at certain times, accompanied by their parents. Nursing professionals collected blood samples (8 mL) in BD Vacutainer™ tubes (Becton, Dickinson and Company, New Jersey, USA). Serum samples were analyzed for the measurement of HDL-c and triglycerides and plasma for glycemia. The semi-automated Bio 200F analyzer (Bioplus, São Paulo, Brazil) was used. Standardized methods quantitatively determined blood variables, following the standards and technical specifications of the reagents used.

The diagnosis of MetS<sup>16</sup> was determined by the presence of at least three of the following alterations: triglycerides  $\geq 100$  mg/dL; HDL-c < 50 mg/dL; glycemia  $\geq 110$  mg/dL; waist circumference  $\geq 75$ th percentile for age and gender; and alteration in BP (diastolic or systolic)  $> 90$ th percentile adjusted for age, height and gender.

Variables were tested for their normality using the Kolmogorov-Smirnov test. Outliers were identified and withdrawn using the interval between quartiles. The Mann-Whitney *U*-test was used to compare independent groups with non-parametric distribution.

For comparison with other studies, the 5th, 10th, 25th, 50th, 75th, 90th, and 95th percentiles were chosen as reference values. Reference curves were created using the Cole LMS method.<sup>17</sup> The LMS method assumes that, for independent data with positive values, the age-specific Box-Cox transformation can be employed to make them have a normal distribution; the *L*, *M*, and *S* values are natural cubic splines with knot sequence without each age range. The sample, in each gender, was separated into age groups with 100 or more individuals, the lowest number considered to be adequate for the LMS method.<sup>17</sup>

The receiver operating characteristic (ROC) curve was used to evaluate the predictive capacity of diagnostic tests. The areas below the ROC curves (AUC) were calculated to assess the discriminatory power of the SFs in the indication of metabolic alterations constituting the MetS. The sensitivity and specificity values of the anthropometric indicators were calculated for each cutoff present in the sample. The cutoff that showed the best equilibrium between sensitivity and specificity was chosen to optimize the association between these two parameters, showing higher accuracy (lower number of false negative and false positive values). The statistical significance of each analysis was verified by the AUC and by the lower limit of the 95% confidence interval  $> 0.5$ .<sup>18</sup>

## Results

A total of 1480 elementary school students from the urban and rural areas of Uberaba, MG, Brazil, with a mean age of 8.55 years (*SD* = 1.53 years), of whom 52.2% were females, participated in the study.

Girls had a higher prevalence of MetS (12.6% vs. 8.5%,  $p < 0.05$ ) and higher SF values in all assessed anatomical points ( $p < 0.05$ ) when compared with boys (Table 1).

Percentile curves (p5, p10, p25, p50, p75, p90, and p95) were created for the four assessed SFs, as well as for their sum, for both genders (Fig. 1). Overall, all SFs showed a linear increase according to age and gender, with higher values being observed in girls.

The four assessed SFs and their sum were accurate in predicting MetS for girls (Table 2) and boys (Table 3). Most of the suggested values are above the 75th percentile for age and gender.

## Discussion

The present study showed percentile SF curves of the central and peripheral regions of the body, as well as the sum of the four SFs to predict the risk of MetS according

**Table 1** Descriptive skinfold characteristics of schoolchildren aged 6–10 years of age, from the municipality of Uberaba, MG, Brazil, by gender.

Variables	Girls (n=773)		Boys (n=707)		p-Value
	Mean (SD)	Median (min-max)	Mean (SD)	Median (min-max)	
BSF (mm)	10.14 (5.15)	9.00 (3.00–29.00)	8.57 (5.29)	6.67 (3.00–28.00)	0.001
TSF (mm)	14.45 (5.66)	13.33 (5.00–35.00)	12.71 (6.67)	10.67 (5.00–35.00)	0.001
SSSF (mm)	11.77 (7.63)	9.00 (4.00–44.00)	9.85 (7.50)	7.00 (4.00–44.00)	0.001
SISF (mm)	14.40 (10.56)	9.67 (3.00–51.00)	11.12 (10.23)	6.67 (3.00–50.00)	0.001
Σ4SFs (mm)	50.93 (28.31)	42.0 (10.00–163.00)	42.37 (29.79)	31.00 (11.00–202.67)	0.001

n, sample number; SD, standard deviation; Min, minimum value; Max, maximum value; TSF, triceps skinfold; BSF, biceps skinfold; SSSF, subscapular skinfold; SISF, suprailiac skinfold; Σ4SFs, sum of the four skinfolds evaluated.

Note: Significant difference between genders  $p \leq 0.05$ , Mann-Whitney test.


**Table 2** Skinfold cutoffs for predicting metabolic syndrome in girls, by age, municipality of Uberaba, MG, Brazil.

Age (years)	Predictors	AUC	p-Value	95% CI	Cutoff (mm)	S (%)	Sp (%)
6	BSF	0.832	0.014	0.709	0.955	11.6	80
	TSF	0.771	0.044	0.566	0.977	15.3	80
	SSSF	0.864	0.007	0.760	0.967	9	100
	SISF	0.805	0.024	0.666	0.943	12.3	80
	Σ4SFs	0.833	0.013	0.708	0.959	40.3	100
7	BSF	0.894	<0.001	0.81	0.977	13.0	88
	TSF	0.927	<0.001	0.874	0.98	17	100
	SSSF	0.924	<0.001	0.871	0.978	13.6	100
	SISF	0.909	<0.001	0.841	0.977	13.6	100
	Σ4SFs	0.927	<0.001	0.872	0.982	57.3	100
8	BSF	0.803	<0.001	0.705	0.902	10	87
	TSF	0.838	<0.001	0.751	0.924	17.6	73
	SSSF	0.874	<0.001	0.775	0.974	14.3	80
	SISF	0.828	<0.001	0.727	0.929	12.3	87
	Σ4SFs	0.858	<0.001	0.770	0.945	57.6	80
9	BSF	0.862	<0.001	0.773	0.951	14.0	86
	TSF	0.844	<0.001	0.731	0.956	17.6	85
	SSSF	0.88	<0.001	0.802	0.959	14	86
	SISF	0.84	<0.001	0.752	0.927	14.3	100
	Σ4SFs	0.871	<0.001	0.788	0.954	66	86
10	BSF	0.764	<0.001	0.627	0.902	14.3	68
	TSF	0.756	<0.001	0.629	0.884	19	75
	SSSF	0.823	<0.001	0.701	0.945	15.6	79
	SISF	0.78	<0.001	0.648	0.911	18.6	80
	Σ4SFs	0.791	<0.001	0.659	0.923	71.6	80

AUC, area under the receiver operating characteristic curve; 95% CI, 95% confidence interval; S, sensitivity; Sp, specificity; TSF, triceps skinfold; BSF, biceps skinfold; SSSF, subscapular skinfold; SISF, suprailiac skinfold; Σ4SFs, sum of the four assessed skinfolds.

to age and gender in Brazilian children. The curves outlined here may be a useful strategy to prevent the risk of MetS in childhood, with the possibility of being used in schools, family health units, clinics, and hospitals. According to the present findings, both the central adiposity, as well as the peripheral adiposity SFs and the sum of four SFs showed cutoffs with high sensitivity and specificity values to predict the risk of MetS in children. Approximately 75% of the children in the present study with SF values above the 75th percentile, regardless of the assessed SF, had MetS.

The prevalence of MetS in Brazilian children and adolescents varied from 0% to 42.4%, according to data from a systematic review.<sup>19</sup> In the present study, MetS was prevalent in 12.6% of the girls and 8.5% of the boys, with a significant difference between the genders. The heterogeneity of definitions and cutoffs for the MetS components may explain, at least partly, the different prevalence rates reported in the literature.<sup>19</sup> However, it is true that the prevalence of MetS has been increasing among children and adolescents, with a significantly higher proportion among those who are obese.<sup>20</sup>



**Figure 1** Percentile curves (p5, p10, p25, p50, p75, p90, and p95) of the triceps, biceps, subscapular, and suprailiac skinfolds, as well as for their sum, of girls and boys aged 6–10 years, Uberaba, MG, Brazil.

The percentile distribution of the triceps and subscapular SFs were performed in 8568 Chinese schoolchildren aged 7–18 years of age<sup>21</sup> and 32,783 North-American children and

adolescents.<sup>10</sup> The percentile values for TSF and SSSF of Chinese and American children were lower than the percentiles of boys and girls in the present study. This result is

**Table 3** Skinfold cutoffs for predicting metabolic syndrome in girls, by age, municipality of Uberaba, MG, Brazil.

Age (years)	Predictors	AUC	p-Value	95% CI	Cutoff (mm)	S (%)	Sp (%)
6	BSF	0.95	0.002	0.902 - 1.00	13.3	100	91.5
	TSF	0.95	0.002	0.893 - 1.00	16.0	100	88.7
	SSSF	0.93	0.004	0.854 - 1.00	8.6	100	82.6
	SISF	0.90	0.007	0.807 - 0.997	11.3	100	81.6
	$\sum 4\text{SFs}$	0.94	0.003	0.87 - 1.00	51.3	100	86.1
7	BSF	0.86	0.007	0.726 - 0.994	8.3	80	72.2
	TSF	0.83	0.015	0.681 - 0.969	11.6	80	72.6
	SSSF	0.90	0.003	0.789 - 1.00	7.6	100	71.6
	SISF	0.82	0.016	0.635 - 1.00	7.0	60	60.7
	$\sum 4\text{SFs}$	0.86	0.007	0.71 - 1.00	32.3	100	66.2
8	BSF	0.84	0.044	0.641 - 1.00	8.3	66.6	63.1
	TSF	0.84	0.048	0.626 - 1.00	11.6	66.6	61.1
	SSSF	0.88	0.025	0.756 - 1.00	9.3	100	73.9
	SISF	0.91	0.016	0.821 - 1.00	13.3	100	82.5
	$\sum 4\text{SFs}$	0.89	0.024	0.761 - 1.00	42.6	100	75.5
9	BSF	0.86	0.001	0.767 - 0.955	10	87.5	68.2
	TSF	0.86	0.001	0.769 - 0.954	14.6	88.8	72.0
	SSSF	0.92	0.001	0.868 - 0.980	12.3	100	83.7
	SISF	0.92	0.001	0.852 - 0.982	14.6	88.8	81.4
	$\sum 4\text{SFs}$	0.90	0.001	0.832 - 0.974	55.3	88.8	81.4
10	BSF	0.80	0.001	0.678 - 0.930	11.3	86.6	70.9
	TSF	0.78	0.001	0.645 - 0.917	17.3	80	79.6
	SSSF	0.77	0.002	0.638 - 0.905	13.3	80	72.3
	SISF	0.79	0.001	0.653 - 0.920	22.0	73.3	86.3
	$\sum 4\text{SFs}$	0.79	0.001	0.661 - 0.920	61.6	80.0	80.3

AUC, area under the receiver operating characteristic curve; 95% CI, 95% confidence interval; S, sensitivity; Sp, specificity; TSF, triceps skinfold; BSF, biceps skinfold; SSSF, subscapular skinfold; SISF, suprailiac skinfold;  $\sum 4\text{SFs}$ , sum of the four assessed skinfolds.

of concern, due to the association between SFs and central obesity, unfavorable lipid profile, increased levels of insulin and BP, and left ventricular mass.<sup>22</sup> The accumulation of subcutaneous fat, translated into high SF values, increase the chances of children having metabolic alterations.

Pre-pubertal Mexican children aged 6–10 years had a three-fold higher chance of having alterations in MetS components when the distribution of SSSF was in the fourth quartile of the sample (low HDL-c levels [OR = 3.16; 95% CI: 1.41–7.10;  $p < 0.01$ ]) and high triglyceride levels (OR = 3.27; 95% CI: 2.02–5.29;  $p < 0.001$ ).<sup>23</sup> However, high values of  $\sum 3\text{SF}$  (TSF + BSF + SSSF; >90th percentile) in German children aged 3–11 years increased the chance of having three or more cardiovascular risk factors by 1.6-fold (95% CI: 1.1–2.2;  $p < 0.05$ ) and the chance of having arterial hypertension by 1.7-fold (95% CI: 1.1–2.7;  $p < 0.05$ ).<sup>24</sup>

The results of the present study demonstrated that all isolated SF measures predict the risk of MetS in both genders. In the scientific investigations related to the subject, studies with TSF and SSSF are predominant.<sup>10,13,21,25</sup> Many studies have used SF values alone as a predictor of the amount of body fat<sup>25–29</sup> and isolated cardiometabolic risk factors.<sup>12,30</sup> Conversely, studies investigating the power of SF in predicting MetS in the pediatric population remain scarce.

The use of the sum of SF absolute values may become an interesting predictor of MetS, as it minimizes biases in predictive body composition equations, as well as

suggesting values that show equilibrium/disequilibrium of body fat distribution (TSF + BSF + SSSF + SISF). Among the different anthropometric measures tested,  $\sum 4\text{SF}$  was the most accurate predictor of MetS in Brazilian girls and boys, with AUC = 0.908 and AUC = 0.897, respectively.<sup>6</sup>

In the present study, the  $\sum 4\text{SF}$  showed an AUC of 0.859 for girls and 0.879 for boys, and the suggested cutoffs showed high sensitivity values (>80%).

To the best of the authors' knowledge, this is the first study with percentile curves of the central and peripheral region SFs, as well as the sum of four SFs, used to predict MetS in a representative sample of Brazilian schools. However, this study had a cross-sectional design, which did not allow the determination of whether alterations in subcutaneous adiposity reflect changes in the MetS components. Thus, longitudinal studies are necessary to define the causal association between MetS and subcutaneous adiposity. It is worth mentioning that, although the study population was delimited among students from the 1st to the 5th year of Elementary School, the State Education Secretariat provided updated data on all Elementary School years, which may have overestimated the sample calculation.

Reference curves from other countries can either underestimate or overestimate the disease due to ethnic, socioeconomic, and cultural differences. The development of percentile curves with populations of different geographic areas is a relevant challenge to allow a more accurate

screening of MetS in young individuals through the evaluation of SFs.

## Funding

Fundação de Amparo à Pesquisa do Estado de Minas Gerais (Fapemig).

## Conflicts of interest

The authors declare no conflicts of interest.

## References

- Mansur AP, Favarato D. Mortality due to cardiovascular diseases in Brazil and in the metropolitan region of São Paulo: a 2011 update. *Arq Bras Cardiol.* 2012;99:755–61.
- World Health Organization. The global burden of disease: 2004 update. Geneva: WHO; 2008.
- Expert panel on detection evaluation, treatment of high blood cholesterol in adults. Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *JAMA.* 2001;285:2486–97.
- Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, et al. Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care.* 2001;24:683–9.
- Srinivasan SR, Myers L, Berenson GS. Predictability of childhood adiposity and insulin for developing insulin resistance syndrome (syndrome X) in young adulthood: the Bogalusa Heart Study. *Diabetes.* 2002;51:204–9.
- Andaki AC, Tinoco AL, Mendes EL, Andaki Junior R, Hills AP, Amorim PR. Anthropometry and physical activity level in the prediction of metabolic syndrome in children. *Public Health Nutr.* 2014;17:2287–94.
- Katzmarzyk PT, Perusse L, Malina RM, Bergeron J, Despres JP, Bouchard C. Stability of indicators of the metabolic syndrome from childhood and adolescence to young adulthood: the Quebec Family Study. *J Clin Epidemiol.* 2001;54:190–5.
- Misra A, Vikram NK, Arya S, Pandey RM, Dhingra V, Chatterjee A, et al. High prevalence of insulin resistance in postpubertal Asian Indian children is associated with adverse truncal body fat patterning, abdominal adiposity and excess body fat. *Int J Obes Relat Metab Disord.* 2004;28:1217–26.
- Ali O, Cerjak D, Kent JW Jr, James R, Blangero J, Zhang Y. Obesity, central adiposity and cardiometabolic risk factors in children and adolescents: a family-based study. *Pediatr Obes.* 2014;9:e58–62.
- Addo OY, Himes JH. Reference curves for triceps and subscapular skinfold thicknesses in US children and adolescents. *Am J Clin Nutr.* 2010;91:635–42.
- Jaworski M, Kulaga Z, Pludowski P, Grajda A, Gurkowska B, Napieralska E, et al. Population-based centile curves for triceps, subscapular, and abdominal skinfold thicknesses in Polish children and adolescents – the OLAF study. *Eur J Pediatr.* 2012;171:1215–21.
- Khadilkar A, Mandlik R, Chiplonkar S, Khadilkar V, Ekbote V, Patwardhan V. Reference centile curves for triceps skinfold thickness for Indian children aged 5 to 17 years and cut offs for predicting risk of childhood hypertension: a multi-centric study. *Indian Pediatr.* 2015;52:675–80.
- World Health Organization. WHO child growth standards: head circumference-for-age, arm circumference-for-age, triceps skinfold-for-age and subscapular skinfold-for-age. Methods and development. Geneva: WHO; 2007.
- World Health Organization. Oral health surveys: basic methods. Geneva: World Health Organization; 1997.
- Sociedade Brasileira de Cardiologia, Sociedade Brasileira de Hipertensão, Sociedade Brasileira de Nefrologia. VI diretrizes brasileiras de hipertensão. *Arq Bras Cardiol.* 2010;95 Suppl. 1:1–51.
- de Ferranti SD, Gauvreau K, Ludwig DS, Neufeld EJ, Newburger JW, Rifai N. Prevalence of the metabolic syndrome in American adolescents: findings from the Third National Health and Nutrition Examination Survey. *Circulation.* 2004;110:2494–7.
- Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stat Med.* 1992;11:1305–19.
- Schisterman EF, Faraggi D, Reiser B, Trevisan M. Statistical inference for the area under the receiver operating characteristic curve in the presence of random measurement error. *Am J Epidemiol.* 2001;154:174–9.
- Tavares L, Yokoo E, Rosa M, Fonseca S. Metabolic syndrome in Brazilian children and adolescents: systematic review. *Cad Saude Colet.* 2010;18:469–76.
- Friend A, Craig L, Turner S. The prevalence of metabolic syndrome in children: a systematic review of the literature. *Metab Syndr Relat Disord.* 2013;11:71–80.
- Ying-Xiu Z, Shu-Rong W. Distribution of skinfold thickness and blood pressure among children and adolescents in Shandong, China. *J Trop Pediatr.* 2011;57:258–62.
- Freedman DS, Serdula MK, Srinivasan SR, Berenson GS. Relation of circumferences and skinfold thicknesses to lipid and insulin concentrations in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr.* 1999;69:308–17.
- Ramirez-Velez R, Suarez-Ortega MF, Aguilar de Plata AC. Association between adiposity and cardiovascular risk factors in prepubertal children. *Endocrinol Nutr.* 2011;58:457–63.
- Haas GM, Liepold E, Schwandt P. Predicting cardiovascular risk factors by different body fat patterns in 3850 German children: the PEP family heart study. *Int J Prev Med.* 2011;2:15–9.
- Kriemler S, Puder J, Zahner L, Roth R, Meyer U, Bedogni G. Estimation of percentage body fat in 6- to 13-year-old children by skinfold thickness, body mass index and waist circumference. *Br J Nutr.* 2010;104:1565–72.
- Freedman DS, Wang J, Ogden CL, Thornton JC, Mei Z, Pierson RN, et al. The prediction of body fatness by BMI and skinfold thicknesses among children and adolescents. *Ann Hum Biol.* 2007;34:183–94.
- Ayatollahi SM, Mostajabi F. Triceps skinfold thickness centile charts in primary school children in Shiraz, Iran. *Arch Iran Med.* 2008;11:210–3.
- Bedogni G, Iughetti L, Ferrari M, Malavolti M, Poli M, Bernasconi S, et al. Sensitivity and specificity of body mass index and skinfold thicknesses in detecting excess adiposity in children aged 8–12 years. *Ann Hum Biol.* 2003;30:132–9.
- Pecoraro P, Guida B, Caroli M, Trio R, Falconi C, Principato S, et al. Body mass index and skinfold thickness versus bioimpedance analysis: fat mass prediction in children. *Acta Diabetol.* 2003;40:S278–81.
- Quadros TM, Gordia AP, Silva RC, Silva LR. Predictive capacity of anthropometric indicators for dyslipidemia screening in children and adolescents. *J Pediatr (Rio J).* 2015;91:455–63.