



ORIGINAL ARTICLE

Role of serum (1,3)- β -D-glucan assay in early diagnosis of invasive fungal infections in a neonatal intensive care unit[☆]



Abd Elazeez Shabaan^{a,*}, Lobna Mahmoud Elbaz^b, Wafaa M. El-Emshaty^c,
Basma Shouman^a

^a Mansoura University, Department of Pediatrics, Mansoura, Egypt

^b Mansoura General Hospital, Department of Pediatrics, Mansoura, Egypt

^c Mansoura University, Department of Clinical Pathology, Mansoura, Egypt

Received 18 April 2017; accepted 18 July 2017

Available online 13 November 2017

KEYWORDS

(1,3)- β -D-glucan;
Invasive candidiasis;
Neonatal sepsis

Abstract

Objectives: To study the microbiological pattern of late onset neonatal sepsis cultures and to assess the diagnostic performance of serum (1,3)- β -D-glucan level for early diagnosis of invasive fungemia in high-risk infants admitted to a neonatal intensive care unit.

Methods: A prospective multicenter clinical trial conducted on infants at high risk for invasive fungal infections, with suspected late onset sepsis, admitted to a neonatal intensive care unit at Mansoura University Children's Hospital and Mansoura General Hospital between March 2014 and February 2016.

Results: A total of 77 newborn infants with high risk of invasive fungal infection were classified based on blood culture into three groups: no fungemia (41 neonates with proven bacterial sepsis), suspected fungemia (25 neonates with negative blood culture), and definite fungemia group (11 neonates with culture-proven *Candida*). The growing organisms were *Klebsiella* spp. (14/54); *Escherichia coli* (12/54); *Staphylococcus* spp. (12/54; coagulase-negative *Staphylococcus* [9/54]; *Staphylococcus aureus* [3/54]); *Pseudomonas aeruginosa* (3/54); and *Proteus* spp. (2/54). Moreover, 11/54 presented *Candida*. Serum (1,3)- β -D-glucan concentration was significantly lower in the no fungemia group when compared with the definite fungemia group. The best cut-off value of (1,3)- β -D-glucan was 99 pg/mL with sensitivity, specificity, positive predictive value, negative predictive value, and accuracy of 63.6%, 95.1%, 77.8%, 90.7%, and 88.5%, respectively.

[☆] Please cite this article as: Shabaan AE, Elbaz LM, El-Emshaty WM, Shouman B. Role of serum (1,3)- β -D-glucan assay in early diagnosis of invasive fungal infections in a neonatal intensive care unit. J Pediatr (Rio J). 2018;94:559–65.

* Corresponding author.

E-mail: ashabaan1973@gmail.com (A.E. Shabaan).

PALAVRAS-CHAVE

(1,3)- β -D-glucano;
Candidíase invasiva;
Sepse neonatal

Conclusion: (1,3)- β -D-glucan assay has a limited sensitivity with excellent specificity and negative predictive value, which allow its use as an aid in exclusion of invasive neonatal fungal infection. Accurate diagnosis and therapeutic decisions should be based on combining (1,3)- β -D-glucan assay with other clinical, radiological, and microbiological findings.

© 2017 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

O papel do (1,3)- β -D-glucano no soro no diagnóstico precoce de infecções fúngicas invasivas em uma unidade de terapia intensiva neonatal

Resumo

Objetivos: Estudar o padrão microbiológico das culturas de sepse neonatal de início tardio e avaliar o desempenho diagnóstico do nível de (1,3)- β -D-glucano no soro para diagnóstico precoce de fungemia invasiva em neonatos de alto risco internados em uma unidade de terapia intensiva neonatal.

Métodos: Ensaio clínico multicêntrico prospectivo conduzido em neonatos internados em uma unidade de terapia intensiva neonatal com suspeita de sepse de início tardio que estavam em risco de infecções fúngicas invasivas no hospital universitário infantil de Almançora e no hospital geral de Almançora entre março de 2014 e fevereiro de 2016.

Resultados: Foram classificados 77 neonatos recém-nascidos com risco de infecção fúngica invasiva, com base na hemocultura, em: grupo sem fungemia, incluindo 41 neonatos com sepse bacteriana comprovada, grupo com suspeita de fungemia, incluindo 25 neonatos com hemocultura negativa; e grupo com fungemia definida, incluindo 11 neonatos com *Candida* comprovada por cultura. Os organismos em crescimento foram: {*Klebsiella* spp 14/54; *E. coli* 12/54; *Staphylococcus* spp 12/54 (*Staph* coagulase negativa 9/54; *Staph aureus* 3/54); *pseudomonas aeruginosa* 3/54 e *Proteus* spp 2/54}, além de 11/54 *Candida*. A concentração de (1,3)- β -D-glucano no soro foi significativamente inferior no grupo sem fungemia em comparação ao grupo com fungemia definida. O melhor valor de corte da (1,3)- β -D-glucano foi 99 pg/mL com sensibilidade, especificidade, valor preditivo positivo, valor preditivo negativo e precisão de 63,6%, 95,1%, 77,8%, 90,7% e 88,5%, respectivamente.

Conclusão: O ensaio de (1,3)- β -D-glucano possui sensibilidade limitada com especificidade e valor preditivo negativo excelentes que possibilitam seu uso e ajudam na exclusão de infecção fúngica invasiva neonatal. O diagnóstico preciso e as decisões terapêuticas devem ter como base a combinação de ensaio de (1,3)- β -D-glucano com outros achados clínicos, radiológicos e microbiológicos.

© 2017 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Neonatal sepsis (NS), together with pneumonia and meningitis, account for 1.4 million annual neonatal deaths worldwide.¹ Newborn infants are at high risk for infection due to underdevelopment of their immune barriers, including fragile skin and relative immune tolerance.² Fungal sepsis is a type of late onset sepsis (LOS) and should be considered in neonates with prolonged hospitalization and in those on prolonged antibiotic treatment.³ The incidence of fungal sepsis ranges from 0.4 to 2 cases per 1000 live births, and from 3.8% to 12.9% among very low birth weight infants.⁴ The most commonly reported risk factors for invasive fungal infection (IFIs) are prematurity, low birth weight, major congenital malformations, exposure to broad spectrum antibiotics, central venous catheters, delayed enteral feeds, prolonged parenteral nutrition, endotracheal intubation, surgery, postnatal steroids, and longer neonatal

intensive care unit (NICU) stay.^{5,6} IFI manifestations are non-specific, and blood culture is considered the gold standard for its diagnosis.⁷ Unfortunately, preliminary results of blood cultures are usually obtained after 48 h or more. Moreover, diagnosis of IFI in neonatal infants is difficult, due to the high rate of false negative in cultures⁸; blood cultures are negative in approximately 50% of cases of autopsy-proven disseminated candidiasis.⁹ Therefore, new tools are required for early diagnosis of IFIs in neonates. 1,3- β -D-Glucan (BG) is a component of the outer cell wall of fungi, including *Candida* spp., *Aspergillus* spp., and *Pneumocystis jiroveci*, and it is released into the blood stream during IFI.¹⁰ To date, only three small trials were conducted in neonates; they concluded that BG is a useful adjunctive diagnostic method of IFI.^{11–13}

The authors hypothesized that early assessment of serum BG in neonates with suspected fungal sepsis is a good substitute to fungal blood culture. Accordingly, this study aimed

to assess the microbiological pattern of neonatal LOS and the diagnostic performance of BG for early diagnosis of IFI in high-risk infants admitted to NICUs.

Subjects and methods

This was a prospective multi-center cohort study conducted in NICUs of the Mansoura University Children's Hospital and of the Mansoura General Hospital from March 2014 to February 2016.

Ethics

The Research Ethics Committee of Mansoura Faculty of Medicine approved the study, and written informed consents were obtained from the parents of all neonates included in the study.

Included subjects

Neonates with clinically suspected LOS (sepsis after 72 h from birth) who were at high risk of IFI were included. Clinical manifestations suggestive of neonatal sepsis were defined in accordance with the Brazilian National Health Surveillance Agency (ANVISA) criteria, in which neonatal sepsis was defined as a systemic response, without any other recognized cause than infection, associated with at least two or more of the following signs and symptoms: thermal instability, apnea, worsening of respiratory discomfort, hemodynamic instability, bradycardia, feeding intolerance, glucose intolerance, hypoactivity, and lethargy.¹⁴

Patients were considered at high risk for IFI if they had three or more of the following criteria: Low birth weight (<2500 g), hospitalization for >3 weeks, prolonged mechanical ventilation (>1 week), systemic antibiotic exposure >72 h, postoperative patients, abdominal wall defects, central venous catheter, arterial catheter >72 h, total parenteral nutrition administration, and persistent severe thrombocytopenia despite second line antibiotics administration.¹³

Excluded subjects

Neonates who received systemic antifungal drugs (prophylactic or therapeutic), intravenous immunoglobulin, albumin, plasma protein, and amoxicillin-clavulanic acid antibiotic were excluded from the study.

All recruited infants underwent LOS investigations: complete blood count, total and differential; quantitative C-reactive protein (CRP); and full septic workup, including blood, urine, CSF cultures, as well as serum BG level. Other laboratory and radiological investigations were performed according to the decision of the attending physician.

Neonates included were categorized according to their blood culture results into three groups. The no fungemia group included 41 neonates with blood culture-proven bacterial sepsis. The definite fungemia group included 11 neonates with blood culture-proven fungemia. Finally, the suspected fungemia group included 25 neonates with negative blood culture for bacteria and fungi, but at high risk of

IFI with abnormal white blood cells (WBCs), low platelets, or elevated CRP.

Laboratory workup

All infants included in the study underwent full septic workup, as well as BG assay, within 24 h of recruitment. A total of 3 mL of blood were collected; 2 mL were used for blood culture using blood culture vials (BD BactecPedsPlus™/F culture vials; Becton Dickinson, Maryland, USA) and the other 2 mL were used for complete blood count (CBC), CRP, and BG analysis.

Methods

Blood culture

An automated continuous-monitoring blood culture system, BD BACTEC™ FX 40 (Becton Dickinson Maryland, USA) was used. Bottles were incubated for five days before being discarded as negative; however, they were sub-cultured according to the laboratory operating procedure if they were flagged as positive before this time. Bacterial isolates were identified through colonies morphology, haemolysis on blood agar, Gram-stained smears, catalase test, coagulase test, and biochemical reactions.

CRP was quantified using latex-enhanced turbid metric assay. The turbidity is measured at 630 nm using HEALES analyzer (Shenzhen Huisong Technology Department Co., Ltd. – China).

BG was detected with the Dynamiker Fungus (Dynamiker Biotechnology, Tianjin, China) in clinical pathology laboratory of Mansoura faculty of medicine. This colorimetric assay based upon a modification of the limulus amebocyte lysate pathway. The absorbance is measured at 405 nm kinetically every 60 s for 40 min at 37 °C with TECAN reader (TECAN, Infinite F 50, Life Sciences, Mannedorf, Switzerland). The concentration of BG is interpreted according to standard curve: values >95, 70–95, and <70 pg/mL were considered positive, equivocal, and negative, respectively, and abnormal results were omitted.

Data analysis

Statistical analysis was done using SPSS software (IBM SPSS Statistics for Windows, version 20.0. NY, USA). The Kolmogorov-Smirnov test was used to check for normal data distribution. Categorical data were described using number and percentage. Continuous data were described using median, range, mean, and standard deviation. The Chi-squared test was used for categorical variables, one-way ANOVA for Gaussian data, and Kruskal–Walls test for non-Gaussian data to compare between three or more groups. The sensitivity and specificity of the BG assay were determined; the positive (PPV) and negative predictive values (NPV) of the test were calculated, and the ROC curve was constructed.




Figure 1 Study flowchart.

Results

During the study period, a total of 77 neonates with LOS at a high risk of IFI were included; the stratification of patients into groups is demonstrated in the study flow chart (Fig. 1). Baseline characteristics of studied groups revealed that gestational age and birth weight were significantly lower in definite fungemia patients when compared with those in the no fungemia and suspected fungemia groups. Postnatal age, Cesarean section delivery, and male gender were significantly higher in definite fungemia patients when compared with no fungemia and suspected fungemia patients (Table 1). Low birth weight, hospitalization >3 weeks, central device, persistent thrombocytopenia, WBC, and CRP were significantly higher in the definite fungemia group when compared with the other two groups, while mechanical ventilation or continuous positive airway pressure (CPAP) >1 week was significantly higher in the definite fungemia group when compared with the no fungemia group only (Table 1). Gram-negative bacteria were more predominant than Gram-positive bacteria, with a frequency of 32/54 and 11/54, respectively. The growing organisms were *Klebsiella* spp. (14/54); *Escherichia coli* (12/54); *Staphylococcus* spp. (12/54; coagulase-negative *Staphylococcus* [9/54]; *Staphylococcus aureus* [3/54]); *Pseudomonas aeruginosa* (3/54); and *Proteus* spp. (2/54). Moreover, 11/54 presented *Candida*. Two neonatal blood cultures revealed mixed growth of *Candida* with bacteria.

The results of the BG assay were negative in 34, equivocal in 23, and positive in 20 cases. The no fungemia group was significantly associated with negative BG results when compared with the definite fungemia group ($p=0.005$); in turn, the definite fungemia group was significantly associated with positive BG results when compared with the no fungemia group ($p=0.001$).

In all studied neonates, the median and ranges of BG concentrations are demonstrated in Fig. 2. BG concentration was significantly higher in the definite fungemia group when compared with the no fungemia and suspected fungemia groups ($p=0.001$ and $p=0.011$, respectively; Fig. 2). The sensitivity and specificity of BG assay at the 99 pg/mL level

were 63.6% and 95.1%, respectively. The PPV and NPV were 77.8% and 90.7%, respectively with accuracy 88.5%. The area under the receiver operating curve (ROC) was 0.837 (95% CI [0.689–0.985]; $p=0.001$; Fig. 3).

Discussion

In spite of the development of new antifungal drugs, mortality and life-threatening complications of IFIs are still frequently reported in critically ill patients.¹⁵ The diagnosis of nosocomial IFI is negatively affected by suboptimal culture results. Therefore, new tools are required for early diagnosis of IFIs in neonates.

In the present study, the microbiological pattern of blood stream neonatal LOS was prospectively tested, and the diagnostic performance of serum BG assay in early diagnosis of IFI in high-risk infants admitted to a NICU was assessed.

In the present study, the most prevalent isolates were Gram-negative bacteria (57.2%), particularly *Klebsiella* spp. accounting for 25.9% of culture positive cases. Similarly, multiple studies reported Gram-negative pathogens as the predominant bacterial isolate in neonatal sepsis, with *Klebsiella pneumoniae* as the most frequently detected organism.^{16,17} Moreover, Al-Shamahy et al. conducted a study in Sanaa (Yemen), and reported a very high rate of Gram-negative bacteria, comprising 97.8% of the total isolates; *Klebsiella* spp. was the most predominant (36.7%) pathogen.¹⁸ In contrast, Hornik et al. found that Gram-positive organisms accounted for 45–75% of causes of LOS in neonates.¹⁹

Fungi represented 20.3% of culture positive cases in the present study. Similarly, a multicenter study conducted by Cotton et al. demonstrated that the rate of IFIs was 2.4–20.4% in extremely low birth weight infants.²⁰ However, a large retrospective study demonstrated a very low incidence of IFIs (0.06%) in neonates >1500 g birth weight.²¹ The high incidence of IFIs in the present study can be explained by the fact that infants who had three or more risk factors for fungal infection were included, as well as by the lack of compliance to strict hygiene in developing countries. In

Table 1 Demographic, clinical, and laboratory characteristics of the studied groups.

	No fungemia <i>n</i> = 41	Suspected fungemia <i>n</i> = 25	Definite fungemia <i>n</i> = 11	<i>p</i> ^a	<i>p</i> ^b	<i>p</i> ^c	<i>p</i> ^d
Gestational age (weeks)	33.41 $\pm$ 3.294	33.08 $\pm$ 3.511	30 $\pm$ 0.256	0.045	0.069	0.037	0.007
Postnatal age (days)	23.76 $\pm$ 8.249	21.08 $\pm$ 7.205	30.91 $\pm$ 6.236	0.003	0.173	0.008	0.001
Birth weight (g)	2100 (850–3900)	1880 (830–4700)	930 (830–3350)	0.038	0.041	0.020	0.018
Male gender	28 (68.3%)	13 (52%)	11 (100%)	0.007	0.187	0.031	0.005
Cesarean section delivery	32 (78%)	19 (76%)	11 (100%)	0.041	0.847	0.047	0.035
Prolonged rupture of membranes	28 (68.3%)	18 (72%)	7 (63.6%)	0.892	0.751	0.770	0.616
Low birth weight (<2500 g)	30 (73.2%)	16 (64%)	10 (90.9%)	0.026	0.432	0.015	0.027
Hospitalization >3 weeks	30 (73.2%)	15 (60%)	11 (100%)	0.038	0.265	0.043	0.016
Mechanical ventilation or CPAP >1 week	36 (87.8%)	22 (88%)	11 (100%)	0.049	1	0.041	0.538
Systemic antibiotic exposure >72 h	33 (80.5%)	20 (80%)	9 (81.8%)	1	1	1	1
Surgical intervention	4 (9.8%)	5 (20%)	0 (0.0%)	0.367	0.282	0.567	0.295
Central venous or arterial catheterization >72 h	23 (56.1%)	15 (60%)	10 (90.9%)	0.003	0.756	0.040	0.016
Total parenteral nutrition administration	34 (82.9%)	20 (80%)	11 (100%)	0.355	0.754	0.322	0.295
Persistent thrombocytopenia ($<100 \times 10^9$ /L)	31 (75.6%)	14 (56%)	10 (90.9%)	0.041	0.047	0.420	0.049
Hemoglobin (g/dL)	11.5 (7.5–18.5)	12.8 (9.3–17.8)	10.3 (9.3–13.5)	0.113	0.947	0.858	0.797
WBCs ($\times 10^9$ /L)	6 (1.9–28.2)	7 (1.9–28.1)	11.5 (3.4–26.3)	0.027	0.077	0.013	0.004
CRP (mg/L)	48 (12–208)	48 (12–112)	96 (12–210)	0.022	0.252	0.024	0.042

Data are expressed as number (percentage), mean $\pm$ SD or median (range).

CPAP, continuous positive airway pressure; WBCs, white blood cells; CRP, C-reactive protein.



^a Comparison between all groups.

^b Comparison between no fungemia and suspected fungemia.

^c Comparison between no fungemia and definite fungemia.

^d Comparison between suspected and definite fungemia.

Chi-square, one-way ANOVA, and Kruskal–Wallis test were used.

**Figure 2** 1,3- β -D-glucan levels in studied neonates.**Figure 3** ROC curve of 1,3- β -D-glucan for discrimination between no fungemia and definite fungemia groups.

the present study, *Candida albicans* was the only isolated fungal growth, which is in agreement with the results of Hope et al., who stated that *C. albicans* is the most frequent *Candida* species causing invasive candidiasis in neonates.²²

Regarding the BG assay in the present study, in the no fungemia group, 23 were found negative, 14 were equivocal, and four were positive. In turn, in the definite fungemia group, seven were found positive, three were equivocal, and one was negative. False-positive BG reactions are suspected to occur in patients treated with intravenous immunoglobulins, albumin, coagulation factors, and plasma protein fraction manufactured by certain fungal vendors, or antibiotics derived from fungal sources as amoxicillin-clavulanic acid, which were all excluded in our study. Patients' exposure to gauze or materials containing glucans during surgery, mucosal damage from chemotherapy or radiotherapy, cross-experimental contamination with BG due to excess manipulation of a sample, certain streptococci, and *P. aeruginosa* may also cause false positive reactivity.^{23,24} Moreover, Zheng et al. reported that post-natal corticosteroids therapy can cause false positive BG result.²⁵ In the present study, two neonates from the no fungemia group had history of recent surgery, with possible exposure to materials that contained glucans. In addition, *P. aeruginosa* was isolated from three neonates within the no fungemia group, which may explain their false positive reactivity. In the suspected fungemia group, nine of the neonates presented positive BG reaction, which might be explained by the low sensitivity of blood culture in the diagnosis of invasive candidiasis, as only 50% of invasive candidiasis are blood-culture positive.⁹ Furthermore, fungi other than *Candida*, such as *Aspergillus*, may be the cause of positive BG in these patients, since blood cultures are almost always negative in disseminated aspergillosis.²⁶ Finally, the false negative result of BG in one patient from the definite fungemia group may be explained by IFI at an early stage.

The sensitivity, specificity, PPV, NPP, and accuracy of the BG assay in the present study at the cut-off value of 95 pg/mL recommended by the manufacturer (after considering the equivocal results to be negative) were 63.6%, 90.2%, 63.6%, 90.2%, and 84.6%, respectively, in comparison to 90%, 56.1%, 35.15, 95.8%, and 63.5%, respectively, at a cut-off value ≥ 70 pg/mL (after considering equivocal results to be positive). At the best obtained cut-off value (99 pg/mL), these levels were 63.6%, 95.1%, 77.8%, 90.7%, and 88.5%, respectively. In the present study, the specificity of BG assay used for early diagnosis of IFIs was excellent (95.1%), while its sensitivity (63.6%) is limited at an optimal cut-off value of 99 pg/mL. The excellent specificity and NPV suggest the good ability of BG assay to exclude IFI in suspected neonates, while its ability to diagnose the disease is limited. Thus, BG assay is better when combined with other clinical, radiological, and microbiological findings to improve its sensitivity. The primary value of the BG assay is to exclude the presence of IFIs based on a negative result, which would eliminate the need for toxic and expensive antifungal therapy.

In a previous study in a similar population of neonates with clinically suspected LOS who were at high risk of fungemia, the BG assay at the 60 pg/mL level showed sensitivity, specificity, PPV, NPV values of 73.2%, 71.0%, 76.9%, and 66.7%, respectively while at the level of 80 pg/mL, these values were 70.7%, 77.4%, 80.6%, and 66.7%, respectively.¹³ In turn, in a retrospective study carried out by Goudjil et al.,

it was reported that BG levels increased in neonatal IFI with an optimal cut-off value for BG positivity >125 pg/mL; at this cut-off value, Sensitivity and specificity were 84% and 75%, respectively.¹¹ Moreover, a study conducted by Zhao et al. found that BG sensitivity in diagnosing IFD at cut-off levels ≥ 10 pg/mL, ≥ 200 pg/mL, and ≥ 400 pg/mL was 68.3%, 49.2%, and 33.3%, respectively while its specificity was 75.6%, 91.3%, and 96.3%, respectively. However, serum BG was measured in their study by a different technique than that of the present study, with BG values <10 pg/mL interpreted as negative and values >20 pg/mL as positive.¹² Furthermore, Liu et al. evaluated the BG assay for diagnosis of candidemia in pediatric patients; they reported results similar to those of the present study, with sensitivity, specificity, PPV, NPV values of 68%, 91%, 66%, and 91%, respectively. They concluded that plasma BG assay has a moderate efficacy to aid the diagnosis of candidemia in pediatric patients, while it is useful to rule out candidemia due to its high NPV.²⁷

In a meta-analysis including 31 studies, it was concluded that BG has a sensitivity of 80% and a specificity of 82% in the diagnosis of IFI. However, most of the studies included in that meta-analysis were conducted in adults.²⁸ In a more recent meta-analysis by Hou et al.,²⁹ in which a total of 1068 patients of different age groups from 11 studies were analyzed, the pooled sensitivity, specificity, positive likelihood ratio, negative likelihood ratio, diagnostic odds ratio, and area under the summary ROC curve, with 95% confidence intervals, were 0.75 (0.63,0.84), 0.87 (0.81,0.92), 5.85 (3.96,8.63), 0.30 (0.20,0.45), 19.53 (11.16,34.18), and 0.89 (0.86,0.91), respectively.²⁹

It is evident that the cut-off value of serum BG can vary according to the kits and techniques used in the assay; in the present study, the best obtained cut-off value for diagnosing IFI was 99 pg/mL with the Dynamiker Fungus technique. Meanwhile, according to a systematic review and meta-analysis conducted by He et al. to evaluate the diagnostic accuracy of serum BG for IFI, the best diagnostic accuracy in the Fungitell assay was observed using a cut-off of 60 pg/mL; in the Fungitec G-Test assay (Seikagaku Corporation, Tokyo, Japan), 20 pg/mL; and in the Wako assay (Wako Chemicals, VA, USA), 11 pg/mL.³⁰

The limitations of the present study include a small sample size, heterogeneous cohort of newborn infants, the fact that two different centers manipulated the samples, and use of blood culture as a gold standard for diagnosis of IFI, which is only positive in 50% of cases. Furthermore, there is the limitation of BG itself as an expensive test, especially for developing countries, and it does not identify species.

It is difficult to measure the actual performance of BG test, due to the poor sensitivity of blood culture. The available data indicates that the BG test has a limited sensitivity and excellent specificity that allow its use as an aid in diagnosis or exclusion of fungal neonatal sepsis. The adjustment of BG cut-off value to 99 pg/mL appears suitable when there are no other causes of false reactivity. The NPV at this level could exclude the presence of fungemia by 90.7% based on a negative result. The authors recommend that accurate diagnosis and therapeutic decisions should be based on combining BG assay with other clinical, radiological, and microbiological findings, due to its limited sensitivity.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Motta M, Zini A, Regazzoli A, Garzoli E, Chirico G, Caimi L, et al. Diagnostic accuracy and prognostic value of the CD64 index in very low birth weight neonates as a marker of early-onset sepsis. *Scand J Infect Dis*. 2014;45:433–9.
- Makhoul IR, Kassir I, Smolkin T, Tamir A, Sujov P. Review of 49 neonates with acquired fungal sepsis: further characterization. *Pediatrics*. 2001;107:61–6.
- Filioti J, Spiroglou K, Panteliadis CP, Roilides E. Invasive candidiasis in pediatric intensive care patients: epidemiology, risk factors, management, and outcome. *Intensive Care Med*. 2007;33:1272–83.
- Schuchat A, Zywicki SS, Dinsmoor MJ, Mercer B, Romaguera J, O'Sullivan MJ, et al. Risk factors and opportunities for prevention of early-onset neonatal sepsis: a multicenter case-control study. *Pediatrics*. 2000;105:21–6.
- Kelly MS, Benjamin DK, Smith PB. The epidemiology and diagnosis of invasive candidiasis among premature infants. *Clin Perinatol*. 2015;42:105–17.
- Benjamin DK Jr, Stoll BJ, Fanaroff AA, McDonald SA, Oh W, Higgins RD, et al. Neonatal candidiasis among extremely low birth weight infants: risk factors, mortality rates, and neurodevelopmental outcomes at 18 to 22 months. *Pediatrics*. 2006;117:84.
- Arnon S, Litmanovitz I. Diagnostic tests in neonatal sepsis. *Curr Opin Infect Dis*. 2008;21:223–7.
- Calley JL, Warris A. Recognition and diagnosis of invasive fungal infections in neonates. *J Infect*. 2017;74:S108–13.
- Clancy CJ, Nguyen MH. Finding the missing 50% of invasive candidiasis: how nonculture diagnostics will improve understanding of disease spectrum and transform patient care. *Clin Infect Dis*. 2013;56:1284–92.
- Kedzierska A, Kochan P, Pietrzyk A, Kedzierska J. Current status of fungal cell wall components in the immunodiagnostics of invasive fungal infections in humans: galactomannan, mannan and (1 $\rightarrow$ 3)-beta-D-glucan antigens. *Eur J Clin Microbiol Infect Dis*. 2007;26:755–66.
- Goudjil S, Kongolo G, Dusol L, Imestouren F, Cornu M, Leke A, et al. (1-3)- β -D-Glucan levels in candidiasis infections in the critically ill neonate. *J Matern Fetal Neonatal Med*. 2013;26:44–8.
- Zhao D, Qiu G, Luo Z, Zhang Y. Platelet parameters and (1, 3)- β -D-glucan as a diagnostic and prognostic marker of invasive fungal disease in preterm infants. *PLOS ONE*. 2015;10:0123907.
- Mackay CA, Ballot DE, Perovic O. Serum 1,3-betaD-glucan assay in the diagnosis of invasive fungal disease in neonates. *Pediatr Rep*. 2011;3:e14.
- De Assis Meireles L, Vieira AA, Costa CR. Evaluation of the neonatal sepsis diagnosis: use of clinical and laboratory parameters as diagnosis factors. *Rev Esc Enferm USP*. 2011;45:33–9.
- De Vlieger G, Lagrou K, Maertens J, Verbeken E, Meersseman W, Van Wijngaerden E. Beta-D-glucan detection as a diagnostic test for invasive aspergillosis in immunocompromised critically ill patients with symptoms of respiratory infection: an autopsy-based study. *J Clin Microbiol*. 2011;49:3783e7.
- Labib AZ, Mahmoud AB, Eissa NA, El Gendy FM, Soliman MA, Aly AA. Early diagnosis of neonatal sepsis: a molecular approach and detection of diagnostic markers versus conventional blood culture. *Int J*. 2013;4:77–85.
- Verma P, Berwal PK, Nagaraj N, Swami S, Jivaji P, Narayan S. Neonatal sepsis: epidemiology, clinical spectrum, recent antimicrobial agents and their antibiotic susceptibility pattern. *Int J Contemp Pediatr*. 2015;2:176–80.
- Al-Shamahy HA, Sabrah AA, Al-Robasi AB, Naser SM. Types of bacteria associated with neonatal sepsis in Al-Thawra University Hospital, Sana'a, Yemen, and their antimicrobial profile. *Sultan Qaboos Univ Med J*. 2012;12:48–54.
- Hornik CP, Fort P, Clark RH, Watt K, Benjamin DK Jr, Smith PB, et al. Early and late onset sepsis in very-low-birth-weight infants from a large group of neonatal intensive care units. *Early Hum Dev*. 2012;88:69–74.
- Cotten CM, McDonald S, Stoll B, Goldberg RN, Poole K, Benjamin DK Jr. The association of third-generation cephalosporin use and invasive candidiasis in extremely low birth-weight infants. *Pediatrics*. 2006;118:717–22.
- Lee JH, Hornik CP, Benjamin DK Jr, Herring AH, Clark RH, Cohen-Wolkowicz M, et al. Risk factors for invasive candidiasis in infants >1500 g birth weight. *Pediatr Infect Dis J*. 2013;32:222–6.
- Hope WW, Castagnola E, Groll AH, Roilides E, Akova M, Arendrup MC, et al. ESCMID* guideline for the diagnosis and management of *Candida* diseases 2012: prevention and management of invasive infections in neonates and children caused by *Candida* spp. *Clin Microbiol Infect*. 2012;18:38–52.
- Pickering JW, Sant HW, Bowles CA, Roberts WL, Woods GL. Evaluation of a (1 $\rightarrow$ 3)-beta-D-glucan assay for diagnosis of invasive fungal infections. *J Clin Microbiol*. 2005;43:5957–62.
- Mennink-Kersten MA, Ruegebrink D, Verweij PE. *Pseudomonas aeruginosa* as a cause of 1,3-beta-D-glucan assay reactivity. *Clin Infect Dis*. 2008;46:1930–1.
- Zheng F, Zha H, Yang D, Deng J, Zhang Z. Diagnostic values and limitations of (1,3)- β -D-glucans and galactomannan assays for invasive fungal infection in patients admitted to pediatric intensive care unit. *Mycopathologia*. 2017;182:331–8.
- Arendrup MC, Fisher BT, Zaoutis TE. Invasive fungal infections in the pediatric and neonatal population: diagnostics and management issues. *Clin Microbiol Infect*. 2009;15:613–24.
- Liu Y, Chen F, Zhu X, Shen L, Zhang SX. Evaluation of a novel plasma (1,3)- β -D-glucan detection assay for diagnosis of candidemia in pediatric patients. *J Clin Microbiol*. 2015;53:3017–20.
- Onishi A, Sugiyama D, Kogata Y, Saegusa J, Sugimoto T, Kawano S, et al. Diagnostic accuracy of serum 1,3- β -D-glucan for *Pneumocystis jirovecii* pneumonia, invasive candidiasis, and invasive aspergillosis: systematic review and meta-analysis. *J Clin Microbiol*. 2012;50:7–15.
- Hou TY, Wang SH, Liang SX, Jiang WX, Luo DD, Huang DH. The screening performance of serum 1,3-beta-D-glucan in patients with invasive fungal diseases: a meta-analysis of prospective cohort studies. *PLOS ONE*. 2015;10:e0131602.
- He S, Hang JP, Zhang L, Wang F, Zhang DC, Gong FH. A systematic review and meta-analysis of diagnostic accuracy of serum 1,3- β -D-glucan for invasive fungal infection: focus on cut off levels. *J Microbiol Immunol Infect*. 2015;48:351–61.