



ORIGINAL ARTICLE

Prevalence, mortality and risk factors associated with very low birth weight preterm infants: an analysis of 33 years^{☆,☆☆}

Julia Damiani Victora^a, Mariangela Freitas Silveira^b, Cristian Tedesco Tonial^{a,*}, Cesar Gomes Victora^b, Fernando Celso Barros^b, Bernardo Lessa Horta^b, Iná Silva dos Santos^b, Diego Garcia Bassani^c, Pedro Celiny R. Garcia^a, Marola Scheeren^a, Humberto H. Fiori^a, Pelotas Cohorts Study Group¹

^a Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), Hospital São Lucas, Programa de Pós-Graduação em Pediatria e Saúde da Criança, Porto Alegre, RS, Brazil

^b Universidade Federal de Pelotas (UFPEL), Programa de Pós-Graduação em Epidemiologia, Pelotas, RS, Brazil

^c University of Toronto, Center for Global Child Health, The Hospital for Sick Children and Department of Pediatrics, Toronto, Canada

Received 26 May 2018; accepted 31 October 2018

KEYWORDS

Preterm;
Very low birth weight;
Cohort studies;
Risk factors;
Prevalence;
Mortality

Abstract

Objective: To assess the prevalence, mortality and risk factors associated with the birth of very low birth weight preterm infants over a period of 33 years.

Methods: Four cross-sectional studies were analyzed, using data from perinatal interviews of birth cohorts in the city of Pelotas collected in 1982, 1993, 2004, and 2015. Based on perinatal questionnaires, anthropometric measurements of newborns and death certificates were analyzed to obtain the prevalence rate, neonatal mortality, and risk factors (maternal age, income and type of delivery) for very low birth weight.

Results: A total of 19,625 newborns were included in the study. In the years 1982, 1993, 2004, and 2015, there were, respectively, 5909, 5232, 4226, and 4258 births. The prevalence of very low birth weight was, respectively, 1.1% ($n=64$), 0.9% ($n=46$), 1.4% ($n=61$), and 1.3% ($n=54$). There was no statistical evidence of an increasing trend over time ($p=0.11$). Among the risk factors, family income in the three poorest quintiles was associated with prevalence rates that were approximately twice as high as in the richest quintile ($p=0.003$). Mortality per 1000 live births for neonates weighing <1500 g decreased from 688 to 259 per thousand from 1982 to 2015 ($p<0.001$), but still represented 61% of neonatal deaths in the latter year.

[☆] Please cite this article as: Victora JD, Silveira MF, Tonial CT, Victora CG, Barros FC, Horta BL, et al. Prevalence, mortality and risk factors associated with very low birth weight preterm infants: an analysis of 33 years. J Pediatr (Rio J). 2018. <https://doi.org/10.1016/j.jpmed.2018.10.011>

^{☆☆} Study conducted at Universidade Federal de Pelotas (UFPEL), Programa de Pós-Graduação em Epidemiologia, Pelotas, RS, Brazil.

* Corresponding author.

E-mail: cristiantonial@gmail.com (C.T. Tonial).

¹ Appendix A lists all the authors of the article.

<https://doi.org/10.1016/j.jpmed.2018.10.011>

0021-7557/© 2018 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALAVRAS-CHAVE

Prematuro;
Muito baixo peso;
Estudos de coorte;
Fator de risco;
Prevalência;
Mortalidade

Conclusion: Although mortality in very low birth weight decreased by more than 60% in recent years, this group still contributes with more than half of neonatal deaths. Low family income remains an important risk factor in this scenario.

© 2018 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Prevalência, mortalidade e fatores de risco associados ao prematuro de muito baixo peso ao nascer: uma análise de 33 anos

Resumo

Objetivo: Verificar a prevalência, mortalidade e fatores de risco associados aos nascimentos de prematuros de muito baixo peso ao nascer (MBPN) ao longo de 33 anos.

Métodos: Série de quatro estudos transversais utilizando dados das entrevistas perinatais das coortes de nascimento da cidade de Pelotas coletados nos anos de 1982, 1993, 2004 e 2015. A partir de questionários perinatais, medidas antropométricas dos recém-nascidos e certidões de óbito, foram analisadas a prevalência, mortalidade neonatal e fatores de risco (idade materna, renda e tipo de parto) para prematuros de muito baixo peso ao nascer.

Resultados: Foram incluídos no estudo um total de 19.625 recém-nascidos. Nos anos 1982, 1993, 2004 e 2015 ocorreram, respectivamente, 5909, 5232, 4226 e 4258 nascimentos. A prevalência de prematuros de muito baixo peso ao nascer naqueles anos foi, respectivamente, de 1,1% (n = 64), 0,9% (n = 46), 1,4% (n = 61) e 1,3% (n = 54). A tendência de aumento durante o período não alcançou significância estatística (p = 0,11). Entre os fatores de risco, a renda familiar nos três quintis mais pobres esteve associada a prevalências cerca de duas vezes mais altas do que no quintil mais rico (p = 0,003). A mortalidade por 1000 nascidos vivos para os neonatos com peso < 1500 g caiu de 688 para 259 por mil ao longo dos anos (p < 0,001), mas ainda representa 61% dos óbitos neonatais em 2015.

Conclusão: Embora a mortalidade nos prematuros de muito baixo peso ao nascer tenha diminuído em mais de 60% nos últimos anos, este grupo ainda contribui com mais da metade dos óbitos neonatais. A baixa renda familiar continua sendo fator de risco importante neste cenário.

© 2018 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In Brazil, since the end of the 1980s, neonatal mortality has been considered the main component of childhood mortality.¹ Preterm birth, with a gestational age of less than 37 weeks, plays an important role in this context, being a potentially preventable cause.²⁻⁴ It is estimated that approximately 12% of the total births of the Brazilian population are preterm, a higher rate than that observed in developed countries.^{5,6}

The causes for preterm birth include factors associated to the mother's age, such as pregnancy in adolescence or above 35 years of age; to the pregnancy, such as a short interval between conceptions, multiple gestations, elective cesarean sections, labor induction, chronic gestational diseases, and infections, as well as socioeconomic and nutritional conditions; and those associated to the fetus, such as genetic diseases.^{5,7}

Neonates with very low birth weight (VLBW) are those whose birth weight is below 1500 g. They constitute a group with high morbidity and mortality, scarcely studied in the literature, and can account for 60% of neonatal deaths.⁸ There is a high correlation between extreme prematurity and VLBW. Based on the Intergrowth-21st study, the 50th

percentile of birth weight for a 32-week-old fetus is equal to 1500 g,⁹ so that, in the absence of accurate gestational age data, temporal trends of VLBW may indicate evolution in extreme prematurity.

In the city of Pelotas, state of Rio Grande do Sul, Brazil, a cohort is started every 11 years, which includes all births that occurred in the city during the year.¹⁰ These cohorts have allowed researchers to assess the evolution of health indicators in the city starting in 1982, 1993, 2004 and, finally, 2015. Based on the data from these studies, the authors attempted to understand the previous and current birth scenario in this group of neonates, in the three decades comprised by the cohorts. Therefore, this study aimed to verify the prevalence, mortality, and risk factors associated with the birth of VLBW neonates throughout this 33-year period.

Methods

Four cross-sectional studies were analyzed using data from the perinatal interviews of the birth cohorts of the city of Pelotas. Data were collected in the years 1982, 1993, 2004 and 2015, and included all live births in the municipality, from the first to the last day of these years. The strategies used for data collection were similar in

Table 1 Birth weight distribution in the birth cohorts of Pelotas, Brazil, in 1982, 1993, 2004, and 2015.

Cohort		Weight groups at birth (g)				Total
		<1500	1500–1999	2000–2499	≥2500	
1982	<i>n</i>	64	109	361	5375	5909
	%	1.1	1.8	6.1	91.0	100
1993	<i>n</i>	46	88	376	4722	5232
	%	0.9	1.7	7.2	90.3	100
2004	<i>n</i>	61	81	281	3803	4226
	%	1.4	1.9	6.6	90.0	100
2015	<i>n</i>	54	94	280	3830	4258
	%	1.3	2.2	6.6	89.9	100
Total	<i>n</i>	225	372	1298	17,730	19,625
	%	1.1	1.9	6.6	90.3	100

Chi-squared test for heterogeneity ($p < 0.001$); chi-squared test for linear trend in VLBW prevalence ($p = 0.11$).

all cohorts: structured questionnaires (perinatal in all cohorts and prenatal in 2015, available at <http://www.epidemiologia-ufpel.org.br/site/content/estudos/index.php>) applied by previously trained interviewers. The mothers were interviewed soon after the delivery and the newborns were measured, weighed, and examined by a team trained and supervised by a pediatrician.^{10,11} Data on the few infants born outside the hospital environment were obtained as soon as the mothers sought care after childbirth. Neonatal mortality (between birth and the 27th day of life, including it) was evaluated through visits to hospitals, cemeteries, notaries, and registries, and were obtained through the SIM (Mortality Information System) from 2004 onwards.^{10–12} The detailed methodological descriptions of each cohort study can be found in the publications by Victora and Barros,¹³ Victora et al.,¹⁴ Santos et al.,¹⁵ and Hallal et al.¹¹

In the present study, the newborns were subdivided according to their birth weight into four subgroups: <1500g, 1500–1999g, 2000–2499g and ≥2500g. The cutoff points of 1500 and 2500g are recommended by the World Health Organization (<http://www.who.int/reproductivehealth/publications/monitoring/9280638327/en/>), and it was decided to subdivide the group between 1500 and 2500g into two subgroups. The distribution of birth weights was described in absolute numbers and percentages. Neonatal mortality was assessed in these groups. For the calculation of gestational age in the 1982 and 1993 cohorts, the date of the last menstrual period reported by the mother was used in the perinatal interview, whereas for the 2004 and 2015 cohorts, the information on gestational age according to the obstetric ultrasound examination was prioritized, and in the absence of this information, the date of the last menstrual period was used. It should also be noted that, in the four cohorts, the gestational age was unknown in approximately 10% of births.

The risk factors studied for VLBW were: monthly family income in minimum wages, maternal age, and type of delivery. The monthly family income was calculated by the sum of the individual incomes and expressed in quintiles of income; maternal age was subdivided into 3 groups: <20 years, 20–34 years, and ≥35 years; and the type of delivery as vaginal or cesarean section. The analyses are shown as tables

and charts. The chi-squared test for proportions was used for the contingency tables.¹⁶ Poisson regression with robust variance was used for raw and adjusted risk factor analyses (expressed in raw and adjusted prevalence for each cohort). Poisson regression was chosen because, in addition to providing correct estimates, it is the most appropriate method for analysis in cross-sectional studies with binary results,¹⁷ making it possible for prevalence ratios to be more easily interpreted and reported when compared with odds ratios calculated by logistic regression.¹⁷ The chi-squared test for linear tendency was used for the analysis of deaths in VLBW over the years. The data were analyzed using the statistical software SPSS (IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY, USA).

This study and all procedures related to the cohorts were approved by the Ethics Committees of the institutions involved in the study. In 1982, the mothers gave verbal consent to participate in the research, whereas from 1993 onwards, their consent was obtained in writing. As of 1996, it has been mandatory to submit projects to the Ethics Committee. In 2004, the study was approved under number 40601116 and, in 2015, under CAAE registration 26746414.5.0000.5313.

Results


A total of 19,625 newborns (NBs) were included in the study. The percentages of newborns (RS) with VLBW ranged from 0.9% to 1.4%, with no evidence of changes over time (chi-squared test for linear trend, p -value=0.11) and are expressed in Table 1. The total number of newborns that were studied in the VLBW group was 225, being assessed as a single group for the analysis of risk factors. The majority of births (89.9%–91.0%) occurred in the group with birth weight ≥2500g in all the years studied. Only nine (5.3%) of the 169 children with VLBW and with known gestational age had 37 or more weeks. In turn, the weight of 99.9% of the full-term children was >1500g.

Table 2 shows that the prevalence rate of VLBW is close to 2.0 in the poorest 60% of cases (corresponding to quintiles 1, 2, and 3 of family income). The main difference is related to the comparison between this group, which includes 60%

Table 2 Risk factors for very low birth weight in the birth cohorts of Pelotas, Brazil, in 1982, 1993, 2004, and 2015.

Prevalence of very low birth weight according to family income, maternal age, and type of delivery. Pelotas, Brazil, 1982–2015.							
Income	% VLBW	Prevalence ratio					
		Raw			Adjusted for the cohort		
		PR (95% CI)	Lower limit	Upper limit	PR (95% CI)	Lower limit	Upper limit
1 st Quintile	1.3	1.68	1.06	2.65	1.67	1.06	2.64
2 nd Quintile	1.6	2.14	1.38	3.31	2.16	1.40	3.33
3 rd Quintile	1.3	1.69	1.07	2.68	1.68	1.06	2.66
4 th Quintile	0.9	1.17	0.72	1.92	1.17	0.71	1.91
5 th Quintile	0.7	1.00			1.00		
<i>p</i> -value		0.004	0.003				
Maternal age							
<20 years	1.4	1.10	0.70	1.72	1.12	0.71	1.76
20–34 years	1.0	0.79	0.54	1.17	0.81	0.55	1.20
≥35 years	1.3	1.00			1.00		
<i>p</i> -value		0.11	0.13				
Type of delivery							
Vaginal	1.1	1.00			1.00		
C-section	1.2	1.14	0.88	1.48	1.07	0.82	1.39
<i>p</i> -value		0.32	0.61				

VLBW, very low birth weight; PR, prevalence ratio.
Statistics used: Poisson regression.

**Figure 1** Neonatal mortality per birth weight groups in the four cohorts.

of the children, and the wealthiest 40% (quintiles 4 and 5), among which the prevalence was below 1%.

The adjustment for the cohort years did not change the observed pattern and the differences between the income groups remained ($p=0.003$). Regarding maternal age, the prevalence of VLBW was slightly lower among mothers aged 20–34, but the difference was not significant ($p=0.13$). No differences were observed between the prevalence of VLBW according to type of delivery ($p=0.61$).

Over the period of 33 years encompassed by the four cohorts, mortality per 1000 live births for neonates weighing <1500 g decreased from 688 in 1982 to 259 in 2015 ($p<0.001$). Among the neonatal deaths, as shown in Fig. 1, the proportion with VLBW increased during the period: 37.9%

in 1982, 43.1% in 1993, 60.0% in 2004, and 60.9% in 2015 ($p<0.001$).

Table 3 shows the results of the multivariate analyses including all factors studied. Only family income showed a significant association ($p=0.003$) with VLBW after adjustment. The prevalence of VLBW was approximately three-fold higher in the poorest 60%, when compared with the wealthiest quintile.

Discussion

The birth cohorts of Pelotas show great potential for the analysis of the temporal evolution of perinatal variables and their risk factors, since they are characterized as population-based studies, covering a broad period of 33 years. The results described herein complement the analyses of the birth weight evolution in Pelotas until 2004, published by Barros et al.¹⁸

The present analyses show a significant decrease in mortality among VLBW newborns throughout the years, whereas the prevalence of births with VLBW has been stable, ranging from 0.9% to 1.4%. A study carried out in the state capital (Porto Alegre) by Da Silva et al., with secondary data, found a VLBW prevalence between 1.1% and 1.3% between 1994 and 2005, with a very slight increase trend ($p=0.049$).

The prevalence rates reported for Porto Alegre are similar to those observed in the present study; it is possible that the borderline level of statistical significance is influenced by the large sample size (more than 20,000 births per year), without indicating a substantial increase in prevalence. The authors report that the increased risk was associated with primiparity.¹⁹

Table 3 Multivariate analysis for very low birth weight in the birth cohorts of Pelotas, Brazil, in 1982, 1993, 2004 and 2015.

Variable	Category	Prevalence ratio	95% CI		p-value%
			Lower	Upper	
Cohort	1982	0.92	0.64	1.32	0.107
	1993	0.73	0.49	1.09	
	2004	1.17	0.81	1.69	
	2015	1.00			
Gender	Male	0.87	0.67	1.14	0.314
	Female	1.00			
Family income	1 st Quintile	1.73	1.08	2.79	0.003
	2 nd Quintile	2.24	1.43	3.50	
	3 rd Quintile	1.74	1.08	2.79	
	4 th Quintile	1.20	0.73	1.98	
	5 th Quintile	1.00			
Type of delivery	Vaginal	1.00			0.173
	C-section	1.21	0.92	1.58	
Maternal age	<20 years	0.98	0.62	1.55	0.270
	20–34 years	0.79	0.54	1.16	
	≥35 years	1.00			

Regarding the risk factors, income was the only studied factor that was associated with VLBW, as observed in the study by Sadovsky et al. for overall prematurity.¹² In the Porto Alegre study, the authors found that the prevalence of VLBW in public hospitals was 67% higher than in private hospitals, which is in agreement with the association observed for socioeconomic status.¹⁹ The same study¹⁹ described an 87% increase in the risk of VLBW for cesarean sections in relation to vaginal deliveries, which was not confirmed in the present study. The association between type of delivery and VLBW is difficult to interpret, since early pregnancy interruptions certainly occur due to medical indication in the case of severe maternal conditions.

Regarding maternal age, a large North American study identified an increased risk for preterm birth (<37 weeks) at the extremes of age, <24 years, and >40 years. Those authors found no association for VLBW preterm births or for gestational age <32 weeks, which is consistent with the present results.²⁰

The significant reduction in neonatal mortality of VLBW newborns over the years can be attributed to improved medical care for this part of the population. The 2015 data showed that this reduction was even more pronounced than in the previous cohorts (Fig. 1).²¹ It should be recalled that, in 1982, there were no neonatal intensive care units in the city, and that the number of beds offered in these units increased from 16 in 1993 to 19 in 2004 and to 29 in 2015. VLBW newborns, however, still account for a large portion of the mortality observed in the neonatal period. In the study "Born in Brazil" of 2011–2012, it was observed that 60% of the neonatal deaths are contained in this VLBW group, which is in agreement with the present findings.⁸

The present study has some limitations. First, only a few risk factors were selected for analysis, based on the literature on VLBW and neonatal mortality determinants. These included family income, type of delivery, maternal age, sex of the newborn, and year of the cohort, with the latter being used to investigate temporal trends. It was not possible to

study public and private hospitals separately, since the analyzed hospitals treat equally patients from the private sector and from the Brazilian Unified Health System.

In conclusion, the prevalence of VLBW remained stable, but there was an evident decrease in mortality among these newborns in recent years. Nonetheless, children with VLBW continue to represent a relevant share of neonatal mortality. The prevention of the birth of very low birth weight children depends not only on the health system improvement but also on policies that can reduce social inequalities.

Institution

Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS)/Postgraduate Program in Pediatrics and Child Health/Hospital São Lucas of PUCRS.

Funding

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

Conflicts of interest

The authors declare no conflicts of interest.

Appendix A. Collaborators: Pelotas Cohorts Study Group

Alicia Matijasevich, Aluísio J.D. Barros, Andréa Damaso Bertoldi, Fernando C. Wehrmeister, Helen Gonçalves, Joseph Murray, Luciana Tovo Rodrigues, Maria Cecília Assumpção, Marlos Rodrigues Domingues, Pedro Rodrigues Curi Hallal

References

1. Victora CG, Aquino EM, do Carmo Leal M, Monteiro CA, Barros FC, Szwarzwald CL. Maternal and child health in Brazil: progress and challenges. *Lancet Lond Engl*. 2011;377:1863–76.
2. Silveira MF, Santos IS, Barros AJ, Matijasevich A, Barros FC, Victora CG. Increase in preterm births in Brazil: review of population-based studies. *Rev Saude Publica*. 2008;42:957–64.
3. Harrison MS, Goldenberg RL. Global burden of prematurity. *Semin Fetal Neonatal Med*. 2016;21:74–9.
4. Santos IS, Matijasevich A, Gorgot LR, Valle NCJ, Menezes AM. Avoidable infant deaths in the 1993 and 2004 Pelotas birth cohorts, Rio Grande do Sul State, Brazil. *Cad Saude Publica*. 2014;30:2331–43.
5. Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller AB, et al. Born too soon: the global epidemiology of 15 million preterm births. *Reprod Health*. 2013;10:S2.
6. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A-B, Narwal R, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet*. 2012;379:2162–72.
7. Frey HA, Klebanoff MA. The epidemiology, etiology, and costs of preterm birth. *Semin Fetal Neonatal Med*. 2016;21:68–73.
8. Lansky S, Friche AA, da Silva AA, Campos D, Bittencourt SD, de Carvalho ML, et al. Pesquisa Nascer no Brasil: perfil da mortalidade neonatal e avaliação da assistência à gestante e ao recém-nascido. *Cad Saude Publica*. 2014;30:S192–207.
9. Villar J, Cheikh Ismail L, Victora CG, Ohuma EO, Bertino E, Altman DG, et al. International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet*. 2014;384:857–68.
10. Barros AJ, Santos IS, Matijasevich A, Araújo CL, Gigante DP, Menezes AM, et al. Methods used in the 1982, 1993, and 2004 birth cohort studies from Pelotas, Rio Grande do Sul State, Brazil, and a description of the socioeconomic conditions of participants' families. *Cad Saude Publica*. 2008;24:S371–80.
11. Hallal PC, Bertoldi AD, Domingues MR, Silveira da MF, Demarco FF, da Silva IC, et al. Cohort Profile: the 2015 Pelotas (Brazil) Birth Cohort Study. *Int J Epidemiol*. 2017.
12. Sadovsky AD, Matijasevich A, Santos IS, Barros FC, Miranda AE, Silveira MF. Socioeconomic inequality in preterm birth in four Brazilian birth cohort studies. *J Pediatr (Rio J)*. 2018;94:15–22.
13. Victora CG, Barros FC. Cohort profile: the 1982 Pelotas (Brazil) Birth Cohort Study. *Int J Epidemiol*. 2006;35:237–42.
14. Victora CG, Hallal PC, Araújo CL, Menezes AM, Wells JC, Barros FC. Cohort profile: the 1993 Pelotas (Brazil) Birth Cohort Study. *Int J Epidemiol*. 2008;37:704–9.
15. Santos IS, Barros AJ, Matijasevich A, Domingues MR, Barros FC, Victora CG. Cohort profile: the 2004 Pelotas (Brazil) Birth Cohort Study. *Int J Epidemiol*. 2011;40:1461–8.
16. Carlin JB, Doyle LW. 5 comparing proportions using the chi-squared test. *J Paediatr Child Health*. 2001;37:392–4.
17. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol*. 2003;3:21.
18. Barros FC, Victora CG, Matijasevich A, Santos IS, Horta BL, Silveira MF, et al. Preterm births, low birth weight, and intrauterine growth restriction in three birth cohorts in Southern Brazil: 1982, 1993 and 2004. *Cad Saude Publica*. 2008;24:S390–8.
19. Da Silva CH, Agranonik M, Da Silva AA, Bettiol H, Barbieri MA, Goldani MZ. Secular trend of very low birth weight rate in Porto Alegre, Southern Brazil. *J Biosoc Sci*. 2010;42:243–53.
20. Fuchs F, Monet B, Ducruet T, Chaillet N, Audibert F. Effect of maternal age on the risk of preterm birth: a large cohort study. *PLoS One*. 2018;13:e0191002.
21. Matijasevich A, Santos IS, Barros AJ, Menezes AM, Albernaz EP, Barros FC, et al. Perinatal mortality in three population-based cohorts from Southern Brazil: trends and differences. *Cad Saude Publica*. 2008;24:S399–408.