

- updated recommendations for a safe infant sleeping environment. *Pediatrics*. 2016;138:e20162940.
28. Psaila K, Foster JP, Pulbrook N, Jeffery HE. Infant pacifiers for reduction in risk of sudden infant death syndrome. *Cochrane Database Syst Rev*. 2017;4:CD011147.
  29. Hauck FR, Thompson JM, Tanabe KO, Moon RY, Vennemann MM. Breastfeeding and reduced risk of sudden infant death syndrome: a meta-analysis. *Pediatrics*. 2011;128:103–10.
  30. Buccini G, Venancio SI. Uso de chupeta em crianças amamentadas: prós e contras. *Sociedade Brasileira de Pediatria*. 2017. Available from: [https://www.sbp.com.br/fileadmin/user\\_upload/Aleitamento-Chupeta\\_em\\_Criancas\\_Amamentadas.pdf](https://www.sbp.com.br/fileadmin/user_upload/Aleitamento-Chupeta_em_Criancas_Amamentadas.pdf) [cited 30.05.19].
  31. Lopes AG, Pereira AC, Fonseca EPd, Mialhe FL. Irregularidades sanitárias na promoção comercial em rótulos de produtos para lactentes e os riscos para a saúde. *Saúde em Debate*. 2017;41:539–52.

32. Sertório SC, Silva IA. As faces simbólica e utilitária da chupeta na visão de mães. *Rev Saude Publica*. 2005;39:156–62.

Gabriela Buccini <sup>a,\*</sup>, Rafael Pérez-Escamilla <sup>a</sup>,  
Sonia I. Venancio <sup>b</sup>

<sup>a</sup> *Yale School of Public Health, New Haven, United States*

<sup>b</sup> *Instituto de Saúde, São Paulo, SP, Brazil*

\*Corresponding author.

E-mail: [gabriela.buccini@yale.edu](mailto:gabriela.buccini@yale.edu) (G. Buccini).

<https://doi.org/10.1016/j.jpmed.2019.06.001>  
0021-7557/

© 2019 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Routine pacifier use in infants: pros and cons<sup>☆</sup>



### Uso rotineiro de chupeta por bebês: prós e contras

Dear Editor,

My Editorial “Routine pacifier use in infants: pros and cons,” recently published in the March/April 2019 issue of *Jornal de Pediatria*,<sup>1</sup> highlighted the conflicting recommendation of the American Academy of Pediatrics (AAP) as opposed to the 10 Steps for Successful Breastfeeding advocated by the World Health Organization/United Nations Children Fund (WHO/UNICEF) regarding pacifier use. As noted, the AAP recommends routine use of pacifiers at sleep time once breastfeeding is established (usually by 3–4 weeks of age)<sup>2</sup> as opposed to the WHO/UNICEF categorical step 9 of the 10 Steps guideline, which stated: “Give no artificial teats or pacifiers to breastfeeding infants.”<sup>3</sup>

It is important to note that, since my writing of the Editorial, the WHO/UNICEF 10 Steps program has been revised; step 9 now states “Counsel mothers on the use and risks of feeding bottles, teats and pacifier”.<sup>4</sup>

While not endorsing the routine use of pacifiers, this still somewhat unbalanced statement clearly retrenches from the previous categorical ban and is more consistent with the consensus (as noted in my Editorial) that the benefits of routine use of pacifiers in reducing the risk of SIDS and in enhancing a calming effect on infant behavior outweigh any theoretical unproven risks.

## Conflicts of interest

The author declares no conflicts of interest.

## References

1. Eidelman AI. Routine pacifier use in infants: pros and cons. *J Pediatr (Rio J)*. 2019;95:121–3.
2. Moon RY. Task force on sudden infant death syndrome, SIDS and other sleep related infant deaths: evidence base for 2016 updated recommendations for a safe infant sleeping environment. *Pediatrics*. 2016;138, pii: e20162940.
3. Saadwh R, Akre J. Ten steps to successful breastfeeding: a summary of the rationale and scientific evidence. *Birth*. 1996;23:154–60.
4. Baby Friendly Hospital Initiative. 10 steps to successful breastfeeding. Available from: <https://www.unicef.org/nutrition/index.breastfeeding-ten-steps.html> [cited 29.05.19].

Arthur I. Eidelman 

*Hebrew University School of Medicine, Shaare Zedek Medical Center, Jerusalem, Israel*

E-mail: [arthur.eidelman@gmail.com](mailto:arthur.eidelman@gmail.com)

<https://doi.org/10.1016/j.jpmed.2019.06.002>  
0021-7557/

© 2019 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

DOIs of original articles:

<https://doi.org/10.1016/j.jpmed.2018.03.002>.

<sup>☆</sup> Please cite this article as: Eidelman AI. Routine pacifier use in infants: pros and cons. *J Pediatr (Rio J)*. 2019;95:621.