

6. Angoorani P, Heshmat R, Ejtahed HS, Motlagh ME, Ziaodini H, Taheri M, et al. Validity of triglyceride–glucose index as an indicator for metabolic syndrome in children and adolescents: the CASPIAN-V study. *Eat Weight Disord.* 2018;23:877–83.

Wendell Arthur Lopes <sup>\*</sup>,  
Gustavo Henrique de Oliveira , João Carlos Locateli ,  
Caroline Ferraz Simões 

*Universidade Estadual de Maringá (UEM), Centro de Ciências da Saúde, Departamento de Educação Física, Curitiba, PR, Brazil*

\*Corresponding author.

*E-mail:* warthurlopes@gmail.com (W.A. Lopes).

<https://doi.org/10.1016/j.jpeds.2019.09.002>

0021-7557/ © 2019 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Answer to the letter ‘‘TyG in insulin resistance prediction’’<sup>☆</sup>



### Resposta à carta ‘‘TyG na predição da resistência à insulina’’

Dear Editor,

We carefully evaluated with great interest the comments made by the authors of the Letter to the Editor and we appreciate the opportunity to answer them.

To calculate the TyG (triglyceride–glucose index) in our study<sup>1</sup>, we used the formula proposed by Simental-Mendía et al.<sup>2</sup> in 2008 and Guerrero-Romero et al.<sup>3</sup> in 2010, described as follows:  $\ln[\text{fasting triglycerides (mg/dL)} \times \text{fasting glucose (mg/dL)}/2]$ . Moreover, the same formula was also used in other studies<sup>4–7</sup> that evaluated the TyG index in predicting insulin resistance in children and adolescents, with similar cutoffs to those identified in our study.

As our main objective was to evaluate the factors associated with the TyG index (dependent variable), linear regression analysis was used (the index was included in the analyses as a quantitative variable). Therefore, regardless of the formula used in the calculation of TyG, there is no change in the observed associations.

Additionally, we redid the TyG index calculation using the following formula:  $\ln[\text{fasting triglycerides (mg/dL)} \times \text{fasting glucose (mg/dL)}]/2$ , which was used by the same authors in another study in 2016<sup>8</sup>. As shown below, the cutoff point identified for the index was lower, as expected. However, the sensitivity and specificity values are similar.

Formula	Cutoff point	Sensitivity (%)	Specificity (%)
$\ln[\text{fasting triglycerides (mg/dL)} \times \text{fasting glucose (mg/dL)}/2]$	7.88	80.0	53.2
$\ln[\text{fasting triglycerides (mg/dL)} \times \text{fasting glucose (mg/dL)}]/2$	4.29	80.0	53.9

<sup>☆</sup> Please cite this article as: Vieira-Ribeiro SA, Fonseca PC, Andreoli CS, Ribeiro AQ, Hermsdorff HH, Pereira PP, et al. Answer to the Letter ‘‘TyG in insulin resistance prediction’’. *J Pediatr (Rio J)*. 2020;96:133–4.

In response to the email sent by a researcher from our institution inquiring about the correct formula for calculating TyG, Fernando Guerrero-Romero points out that: ‘‘Both procedures appropriately identify the presence of insulin resistance; however, using the procedure  $[\ln(\text{glucose} \times \text{triglycerides}/2)]$  gives higher cutoff values for the TyG.’’

We believe in the importance of the standardized formula use for the index calculation; however, to the best of our knowledge, there is no opinion from the authors who validated it regarding which one would be better.

Finally, we would like to highlight the importance of the observations made by the authors of the Letter, as they will greatly contribute to other studies that adopt the TyG index.

### Conflicts of interest


The authors declare no conflicts of interest.

### References

- Vieira-Ribeiro SA, Fonseca PC, Andreoli CS, Ribeiro AQ, Hermsdorff HH, Pereira PF, et al. The TyG index cutoff point and its association with body adiposity and lifestyle in children. *J Pediatr (Rio J)*. 2019;95:217–23.
- Simental-Mendía LE, Rodríguez-Moraan M, Guerrero-Romero F. The product of fasting glucose and triglycerides as surrogate for identifying insulin resistance in apparently healthy subjects. *Metab Syndr Relat Disord*. 2008;6:299–304.
- Guerrero-Romero F, Simental-Mendía LE, González-Ortiz M, Martínez-Abundis E, Ramos-Zavala MG, Hernández-González SO, et al. The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp. *J Clin Endocrinol Metab*. 2010;95:3347–51.
- Kim JH, Park SH, Kim Y, Im M, Han HS. The cutoff values of indirect indices for measuring insulin resistance for metabolic syndrome in Korean children and adolescents. *Ann Pediatr Endocrinol Metab*. 2016;21:143–8.
- Kang B, Yang Y, Lee EY, Yang HK, Kim HS, Lim SY, et al. Triglycerides/glucose index is a useful surrogate marker of insulin resistance among adolescents. *Int J Obes (Lond)*. 2017;41:789–92.
- Moon S, Park JS, Ahn Y. The cut-off values of triglycerides and glucose index for metabolic syndrome in American and Korean adolescents. *J Korean Med Sci*. 2017;32:427–33.
- Angoorani P, Heshmat R, Ejtahed HS, Motlagh ME, Ziaodini H, Taheri M, et al. Validity of triglyceride–glucose index as an

indicator for metabolic syndrome in children and adolescents: the CASPIAN-V study. *Eat Weight Disord.* 2018;23:877–83.

8. Guerrero-Romero F, Villalobos-Molina R, Jiménez-Flores, Simental-Mendía LE, Méndez-Cruz R, Murguía-Romero M, et al. Fasting triglycerides and glucose index as a diagnostic test for insulin resistance in young adults. *Arch Med Res.* 2016;47:382–7.

Sarah A. Vieira-Ribeiro <sup>a,\*</sup>, Poliana C.A. Fonseca<sup>a</sup>,  
Cristiana S. Andreoli<sup>a</sup>, Andréia Q. Ribeiro<sup>b</sup>,  
Helen H.M. Hermsdorff<sup>b</sup>, Patrícia F. Pereira<sup>b</sup>,  
Silvia E. Priore<sup>b</sup>, Sylvia C.C. Franceschini<sup>b</sup>

<sup>a</sup> *Universidade Federal de Viçosa (UFV), Viçosa, MG, Brazil*

<sup>b</sup> *Universidade Federal de Viçosa (UFV), Departamento de Nutrição e Saúde, Viçosa, MG, Brazil*

\* Corresponding author.

*E-mail:* [sarah.vieira@ufv.br](mailto:sarah.vieira@ufv.br) (S.A. Vieira-Ribeiro).

<https://doi.org/10.1016/j.jped.2019.09.003>

0021-7557/ © 2019 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).