



REVIEW ARTICLE

Allergic rhinitis and sleep disorders in children – coexistence and reciprocal interactions



Cláudio D’Elia ^{a,b,*}, David Gozal ^c, Oliviero Bruni ^d, Ekaterini Goudouris ^e, Miguel Meira e Cruz ^{b,f}

^a Hospital Lusíadas, Departamento de Pediatria, Lisbon, Portugal

^b Centro Europeu do Sono, Portugal

^c The University of Missouri School of Medicine, Child Health Research Institute, Department of Child Health, Columbia, United States

^d La Sapienza University, Department of Social Development and Psychology, Rome, Italy

^e Universidade Federal do Rio de Janeiro, Instituto Martagão Gesteira de Puericultura e Pediatria (IPPMG), Faculdade de Medicina, Departamento de Pediatria, Rio de Janeiro, RJ, Brazil

^f Lisbon School of Medicine, Sleep Unit, Centro Cardiovascular da Universidade de Lisboa, Lisbon, Portugal

Received 21 July 2021; accepted 11 November 2021

Available online 31 December 2021

KEYWORDS

Allergic rhinitis;
Sleep disorders;
Sleep-disordered breathing;
Obstructive sleep apnea

Abstract

Objective: To review, critically analyze and synthesize knowledge from the international literature regarding the association between allergic rhinitis (AR) and sleep disorders, the impact of AR treatment on children’s sleep, and lay the foundation for future research on this topic.

Source of data: A literature search using PubMed database including original and review articles, systematic reviews and meta-analyses using keywords related to AR, sleep disorders and sleep-disordered breathing.

Synthesis of data: Sleep is fundamental to health, and its assessment and control of conditions that trigger or aggravate disturbances are of the uttermost importance. Allergic rhinitis (AR) is common in children and may interfere with both their quality of life and quality of sleep. It has emerged as one of the most important risk factors for habitual snoring in children and appeared to increase the risk of Obstructive Sleep Apnea (OSA), with AR severity exhibiting a significant and independent association with pediatric OSA severity. However, in some studies, those associations between AR and OSA in children are not very consistent.

Conclusions: A substantial level of controversy exists regarding the interactions between AR and OSA in children. Notwithstanding, identifying and treating AR in clinical settings is probably an important step toward improving symptoms and preventing deterioration of sleep quality in children and may improve the severity of underlying OSA. Considering the high prevalence, morbidity, economic and social implications of both AR and sleep problems, it is crucial that healthcare

* Corresponding author.

E-mail: claudio.elia@lusiadas.pt (C. D’Elia).

providers improve their understanding of the relationships between those conditions among children.

© 2021 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The mechanisms, signs, and symptoms of allergic conditions such as AR present with clear circadian and ultradian oscillations with a striking day-night pattern. Patients with seasonal and perennial AR complain of disturbed sleep at night, as well as symptoms of awakening in the morning.¹ The earliest chronobiological characterization of AR was by Trousseau in 1865. Since then, many studies have examined the day-night variation and intensity of AR,² with several controversial issues remaining under debate.

AR classical presentation involves sneezing, pruritus, rhinorrhea, and nasal congestion, as a conglomerate of symptoms and functional changes either predisposing to or worsening underlying sleep disorders, including sleep apnea.

OSA is a relatively prevalent condition in children, affecting 1%–5% of children aged 2–8 years, and is caused or aggravated by a variety of different conditions, including AR.³

Despite the low quality of evidence from many studies examining the associations of AR with sleep impairment, both in children and adults, it has been shown that AR patients present with higher levels of disturbance and sleep latency scores and a decreased sleep efficiency score using Polysomnography (PSG). AR was also associated with a higher risk of nocturnal sleep-related dysfunctions, including insomnia, nocturnal enuresis, restless sleep, sleep-disordered breathing (SDB) as OSA or snoring. Additionally, AR was found to be associated with a higher risk of daytime sleep-related impairment.⁴

The management of children with chronic disorders such as AR has become a frequent component of the activities within any pediatric clinic and usually involves both the general pediatricians and pediatric subspecialists. Because of the high prevalence of AR and OSA, its associated morbidity, and the economic and social implications of either respiratory allergies or sleep problems, it is imperative that healthcare providers acquire a deeper understanding of the relationships between both conditions in children and their mutual interaction.

In light of the contradictory findings on the relationship between AR and sleep and its disorders, the authors aimed to review a potential pathophysiological matrix for the crosstalk between AR and sleep disorders, as well as the implications of rhinitis control measures on sleep.

Allergic rhinitis prevalence in children

Allergic rhinitis (AR) is a widespread and common disease. In the United States, the Agency for Healthcare Research and Quality estimates the prevalence of AR in adults to be between 10% and 30%, and several studies have estimated the prevalence of AR in children to be closer to 40%.^{5,6} In the International Study of Asthma and Allergies in Childhood

(ISAAC), the overall prevalence of rhinoconjunctivitis was 8.5% in children aged 6–7 years and up to 14.6% in 13–14-year-old children.⁷ AR is a major non-communicable disease that can impose a significant impact on quality of life, considering its effects during both wakefulness and sleep.⁸

Although the prevalence of AR is clearly heterogeneous and greatly varies with location and socioeconomic status in different countries, the apparent escalation in AR prevalence in children over the last several decades has been paralleled by a similar increase in the frequency of other allergy-related disorders, such as asthma and atopic eczema.⁹ AR can present at any age but will usually become manifest after 2 years of age, with incremental increases in prevalence in older children and adolescents.⁹

Basic mechanisms of allergy

As pathophysiological pathways of allergic disorders are well studied, it is known that after allergen sensitization, whereby antigen is presented to naïve T lymphocytes via an antigen-presenting cell (APC), T cells will undergo differentiation into Th2 cells upon stimulation by thymic stromal lymphopoietin (TSLP) and other cytokines (IL-4, IL-5, IL-13).

Th2 lymphocytes bind to B cells via major histocompatibility complex class II (MHCII) and secrete IL-4 and IL-13. These cytokines induce the B cell to undergo class switching and to produce IgE specific to the sensitizing antigen.¹⁰ Besides stimulating IgE production, Th2 cells produce a variety of cytokines (type-2 cytokines), which enhance eosinophilopoiesis (IL-5), mast cell development (IL-9), and goblet cell hyperplasia.

Type I hypersensitivity processes involve a 2-step temporal process: the early response, reaching a maximum within 30 minutes and usually resolving within 1–3 hrs, and the late response, whereby symptoms start to increase again after 3–4 hrs and reach their peak over 6–12 hrs. The allergen-induced responses are initiated when IgE binds to high-affinity receptors (FcεRI) located on the surface of mast cells and basophils. IgE is subsequently cross-linked by allergen, causing mast cell degranulation, and the release of preformed mediators such as histamine, neutral proteases, and chemotactic factors, along with the activation of eicosanoid pathways to produce newly formed mediators, such as cysteinyl leukotrienes (CysLTC4 and CysLTD4) and their metabolite cysLTE4, as well as prostaglandins (PG), especially PGD2. CysLTs are the mediators underlying nasal congestion and nasal mucosal swelling by increasing microvascular permeability and stimulating mucus secretion. Histamine also contributes to inflammation during the early phase response,¹¹ whereas in the late response, these early mediators are involved in further recruitment of cells including eosinophils, basophils, monocytes, macrophages, and lymphocytes into the inflammatory milieu.¹²

Diurnal variation in AR symptoms

AR and other allergic diseases are underpinned by frank day-night oscillatory patterns either in clinical symptoms, laboratory parameters, or response to treatment.¹³ In most AR patients, symptoms worsen overnight or early in the morning, thereby compromising night-time sleep and resulting in poor daytime quality of life.^{2,13-15} The circadian nature of allergic diseases pathophysiology is well documented, and evidence has been accumulated that the circadian clock regulates IgE/mast cell-mediated allergic reactions.¹³ It was demonstrated an occurrence of 24h clock based variations in histamine release, cysteinyl leukotriene production, and IL-6/IL-13 expression in mast cells following IgE-mediated activation, eventually due to circadian variation in FcεR1a expression.¹⁶ Although rhythmic humoral signals such as cortisol from the adrenal gland¹⁷ and periodically recurrent activity of the autonomic nerves¹³ have been implicated in diurnal oscillations, the biological basis of this phenomenon remains poorly understood.¹³

It was demonstrated that endogenous glucocorticoids could synchronize circadian expression of the clock genes, and the core circadian clock gene *Per2* may contribute to symptomatic temporal variations in AR by its possible anti-inflammatory effect on controlling rhythmic changes in levels of Th2- and Th17-associated transcription factors (eg, GATA-binding protein 3 [GATA3], and ROR γ t).¹⁸ Recent reviews highlighted that oscillatory allergic reactions are generated by rhythmic expression of key molecules in the pathophysiology controlled by the circadian clock.¹⁹

The worsening of symptoms during nighttime may be related to exacerbation of inflammatory processes, usually as a manifestation of the late-phase allergic response to provoking environmental antigens imposed earlier in the day, and such processes also involve key neuroendocrine rhythmic dynamics.²⁰ The day-night differences in symptom intensity, however, may also represent, at least to some extent, circadian rhythm differences in vulnerability to antigen exposure as diurnal variation in nasal reactivity suggests a circadian rhythm-dependent difference in the ability of specific antigens as well as non-specific chemical substances to induce pro-inflammatory activities.²¹

When considering perennial AR, it is suspected that greater exposure loads to mites and their associated antigens at night, when in contact with mattress and pillows, clearly account for a substantial proportion of the nocturnal symptomatic severity increases²⁰ with effects on sleep.²²

Sleep disorders and its consequences

Sleep is an important pillar of a healthy life. Yet, sleep disorders are common in children and adolescents, with an estimated prevalence ranging from 11% to 47%.²³ OSA, as a major sub-group of overall sleep disorders, may conservatively affect up to 4% of children.^{3,24} Impaired nocturnal sleep in children results in daytime fatigue and sleepiness, often associated with altered immune function, anxiety, attention deficit problems, memory deficits, behavioral issues, irritability, depression, growth retardation, hormone imbalance, hypertension, poor academic performance, increased accident rates, substance abuse, as well as

increased risk for cardiovascular and metabolic dysfunction, ultimately leading to lower overall and health-related quality of life.²⁵⁻²⁷ As such, any condition as AR that affects sleep integrity and continuity and reduces its restorative properties is liable of generating symptoms that are not directly related to the disease per se but rather reflect the consequences of disrupted sleep homeostasis.

Association between AR and SDB

In a systematic review analyzing the association between AR and SDB in children, it was concluded that although the majority of the studies showed a significant association between AR and SDB, in all of these studies, the level of evidence was 3b and 4, for an overall grade of B—evidence (Oxford Evidence-Based Medicine Center). As such, the evidence for the association between rhinitis and OSA in children is not very strong, in part because most pediatric studies, until not long ago, used indirect measurements to document SDB (i.e., snoring) instead of overnight PSG to confirm the diagnosis of pediatric OSA.²⁸ Yet, significant associations between snoring and AR were reported in pre-school and primary school children^{29,30} and in school children.³¹

SDB is an important component of the morbidity associated with AR, and it is currently accepted that AR increases the risk of OSA.^{32,33} Although AR severity was found to have a significant and independent association with the severity of OSA in children,³⁴ it was not confirmed in other PSG-diagnosed studies, despite the high prevalence of AR in patients with SDB.³⁵⁻³⁸ The Apnea-hypopnea index (AHI) during rapid eye movement (REM) sleep in children with moderate-severe OSA was significantly increased in subjects with rhinitis and OSA compared with those with OSA alone.³⁵ The AHI severity (determined by total events occurring in REM + NREM) may be affected by different levels of nasal congestion occurring in REM sleep versus NREM sleep. Given that REM sleep occupies only a small proportion of the night (around 20-25%), breathing abnormalities during REM sleep may not necessarily correlate with the overall AHI severity score (which includes NREM sleep that represents 75- 80% of the night).³⁵

Although controversial issues subsist regarding the role of nasal inflammation in primary snoring and OSA, allergic rhinitis appears to be present in 35% of children with primary snoring and in 6% with OSA. On the other hand, non-allergic rhinitis was found in 19% of children with primary snoring and in 34% with OSAS. In this sense, nasal cytology could provide interesting information that can be used to plan possible treatment strategies.³⁹

Accordingly, rhinitis may be considered as a set of symptoms potentiating SDB, whereas in other studies, symptoms of AR may be considered an independent predictor for SDB and failure of adenotonsillectomy (AT) in children.²⁵

Notwithstanding, a recent study that enrolled 1.578 children with habitual snoring did not find a significant association between AR and OSA,⁴⁰ and no association was found between AR and OSA severity in children who were admitted for inpatient AT.⁴¹ Therefore, the presence of enlarged tonsils and adenoids as a dominant factor in the genesis and severity of OSA in children might mask the relative contributions of AR to this condition considering the dominant

contribution of ATH. The authors should, however, consider that before ATH reaches a stage that results in OSA, the proliferation of such lymphoid tissues may be driven by underlying AR, such that the contribution of AR to upper airway dysfunction may be more readily apparent during the early pathophysiological stages of OSA, and progressively lose its prominence as ATH and OSA become more firmly established.

Atopic status and SDB

In one cohort of 105 young children with habitual snoring, significant relationships emerged between this condition and atopic status.⁴²

In a retrospective study with 120 children treated for snoring, the percentage of children with abnormal serum levels of total IgE or abnormal allergen-specific IgEs was comparable in PSG- diagnosed OSA and non-OSA groups. The proportions of abnormal total airway resistance and of abnormal serum levels of total IgE were significantly higher in children with severe OSA compared with children with mild or moderate OSA. The authors suggested that allergic sensitization might contribute to the exacerbation of existing OSA in children.⁴³

Association between AR and sleep disorders

It is worth pointing out that AR may cause daytime fatigue or sleep fragmentation without having evident effects on breathing.⁴⁴ Over the last 15 years, about one-third of the studies failed to show the presence of a significant association between rhinitis and overall sleep disorders (Table 1).^{35,38,40,41,45-63} The very low-grade level of evidence was further corroborated in a more recent systematic review and meta-analysis evaluating the association between AR and sleep, based on observational investigations in children and adults.⁴ Adjusted odds ratios revealed that AR was associated with a higher risk of nocturnal dysfunction, including insomnia, nocturnal enuresis, restless sleep, SDB, OSA, snoring, daytime dysfunction, including difficulty waking up, daytime sleepiness, morning headache, and the use of sleep medications.⁴

The authors surmise that improved delineation of the criteria for AR definition and use of PSG and other objective criteria for sleep assessments will be required in future prospective studies to more compellingly demonstrate or refute the impact of AR on sleep and vice-versa.

It was demonstrated in a few PSG studies in children that the sleep architecture may be altered by AR. The percentage of time spent in the REM sleep stage was lower among children with AR and SDB, without sleep apnea.^{37,38}

AR and sleep questionnaires

Each sleep disturbance detection method has its own reliability and precision as well as its particular indication according to the intended objective. Among the different available tools, the most commonly used for screening sleep quality, sleep habits and disorders, are validated

questionnaires. This option has a very low cost, is easy to administer, and does not require a sleep center. Furthermore, it could have a relevant role as an adjuvant for diagnostic assessment when polysomnography is not available. Another advantage of the use of questionnaires is the allowance of self-administration with high accuracy, as shown in different studies.⁶⁴

Among the different surveys and questionnaire-based tools investigating sleep habits in children are the Children's Sleep Habits Questionnaire (CSHQ). This is a retrospective parent-report questionnaire that was developed in the United States to evaluate sleep behavior in children. The questions were selected to include the symptom presentations of the most common pediatric sleep disorders, according to the International Classification of Sleep Disorders. The Portuguese version of the CSHQ (CSHQ-PT) showed psychometric properties that were comparable to versions from other countries and adequacy for the screening of sleep disturbances in children from 2 to 10 years old.⁶⁵

A case-control study reported that children with AR have higher scores on the CSHQ than the scores of healthy children, indicating a greater probability of sleep disorders being present.⁴⁶ Using the previously reported CSHQ score cutoff, 83% of the children with persistent moderate-to-severe AR would be classified as having sleep disorders, with significantly higher CSHQ scores being found among those children with uncontrolled AR.⁴⁶

Chronic rhinitis has also been associated with habitual snoring in a questionnaire-based survey in Greece on 3,680 children, in whom habitual snoring was present in 5.3%, 4%, and 3.8% in the children of 1-6, 7-12, and 13-18-year age-groups, respectively.⁶⁶

Interactions between allergic rhinitis and sleep problems

AR can affect sleep via several mechanisms (Figure 1). In light of the increased nasal airflow resistance, it is not surprising that AR is considered a risk factor for OSA. In addition, AR can foster enlargement of tonsils and adenoids due to the chronic inflammation in the nasal cavity and nasopharynx and can also promote the evolution of an elongated face due to oral breathing rather than nasal breathing. Taken together, all these factors result in a higher risk for the presence of a smaller and more collapsible upper airway.³²

ATH is by far the major pathophysiological contributor to SDB in children, and its frequency is higher in children with the allergic disease compared to controls,⁶⁷ and allergic disease is also considered a risk factor for adenotonsillar regrowth in children after AT.⁶⁸

Worsening of AR during sleep may partially relate to the typical recumbent position. This position can worsen nasal obstruction due to an increase in nasal turbinate edema.⁶⁹

The nose, when obstructed, can act as an opposing force to favorable upper airway airflow. Inspiration generates negative intrathoracic pressures compared with atmospheric pressure, thereby sucking air into the lung cavity. Nasal obstruction secondary to nasal congestion influences this

Table 1 Summary of published studies examining the relationship between AR and Sleep Disorders in children and adolescents over the last 15 years.

Authors, year, reference	Age	Sample size/AR cases	Study type	Measurement-AR	Measurement sleep outcome	Outcomes
2020 Xu Z et al. ⁴⁰	2-15	1578/244	Cross-sectional	Face-to-face interview	PSG	OSA
2018 Shen L et al. ⁴⁵	<14	676/121	Case- controlled	Self-reported Rhinolaryngosc opy	PSG	OSA
2018 Loekmanwidjaja J et al. ⁴⁶	4–10	167/112	Case- controlled	Medical records	Self-reported	Parasomnias, daytime sleepiness
2018 Lai PH et al. ⁴⁷	<18	655.529/327.928	Cohort	Medical records	Medical records	OSA, nocturnal enuresis
2018 Bilgilişoy Filiz et al. ⁴⁸	8–18	287/143	Case- controlled	Medical records, clinical examination and SPT	Face-to-face interview	Sleep quality (PSQI outcomes)
2017 Tsai JD et al. ⁴⁹	5–18	8616/4191	Case- controlled	Medical records, clinical examination	Medical records	Nocturnal enuresis
2017 Nguyen- Hoang Y et al. ⁵⁰	6-17	85/52	Case- controlled	Clinical examination	Medical records	OSA in asthmatics
2016 Di Francesco RC, Alvarez J ³⁸	3–14	135/57	Case- controlled	Self-reported, Rhinoscopy and SPT	PSG	PSG outcomes (AHI)
2015 Poachanukoon O, Kitcharoensakkul M ⁵¹	6-15	175/65	Case- controlled	Medical records, clinical examination and SPT	Self-reported	Sleep duration, difficult waking up, snoring, morning headache, mouth breathing, night sweating, nocturnal enuresis, OSA daytime sleepiness, restless sleep, SDB, sleep bruxism
2014 Weinstock TG et al. ⁴¹	5-9.9	464/110	Cross-sectional	Physician-diagnosed	PSG	AHI severity
2014 Ng DK et al. ⁵²	15.3 ± 1.7	175/65	Cohort	Self-reported	Self-reported	Snoring
2014 Huseni S et al. ³⁵	2-13	145/63	Cross-sectional	Medical records	PSG	AHI during REM and non-REM
2012 Fadzil Abdullah AA et al. ⁵³	6-10	550/22	Case-controlled	survey booklet	survey booklet	SDB
2012 So HK et al. ⁵⁴	7–10	6381/2661	Cross-sectional	Self-reported	Self-reported	Night sweating
2012 Ishman SL et al. ⁵⁵	0-18	134/21	Case-controlled	SPT or RAST	PSQ; OSA18; PDSS	SDB, sleepiness
2010 Bhattacharjee R et al. ⁵⁶	6.9±3.8	578/212	Case- controlled	Medical records	PSG	Residual OSA post-AT
2010 Urschitz MS et al. ⁵⁷	7-12	1144/712	Cross-sectional	Physician-diagnosed	HPO Questionnaire	OSA
2010 Li S et al. ⁵⁸	5–14	6369/2823	Cross-sectional	Self-reported	Self-reported	Snoring
2009 Sogut A et al. ⁵⁹	12-17	1030/36	Case controlled	Questionnaire	Questionnaire	Snoring
2008 Liukkonen K et al. ⁶⁰	1-6	1471/236	Case controlled	Self-reported	SDSC	Snoring
2008 Petry C, et al. ⁶¹	9-14	998/299	Cross-sectional	Questionnaire	Questionnaire	SDB
2005 Sogut A et al. ⁶²	3-11	1198/332	Case controlled	Self-reported	Self-reported	Snoring
2005 Ng DK et al. ⁶³	6–12	3047/1242	Cross-sectional	Self-reported	Self-reported	Daytime sleepiness, OSA

AHI, Obstructive apnea–hypopnea index; AR, allergic rhinitis; ARIA, Allergic Rhinitis and its Impact on Asthma; AT, adenotonsillectomy; ESPRINT-15, AR health-related quality of life questionnaire for children; ESS, Epworth Sleepiness Scale; HPO, Home pulse oximetry; OSA 18, Obstructive Sleep Apnea-18 survey; OSA, obstructive sleep apnea; PDSS, Pediatric Daytime Sleepiness Scale; PSG, polysomnography; PSQ, Pediatric Sleep Questionnaire; PSQI, Pittsburgh Sleep Quality Index; RAST, in vitro allergy testing; REM, rapid eye movement; RLS, restless legs syndrome; SDB, sleep-disordered breathing; SDSC, Sleep Disturbance Scale for Children; SPT, allergy skin-prick testing.




Figure 1 Rhinitis and Sleep interactions in children. UAW, Upper airways; ROS, Reactive oxygen species.

pressure differential by limiting airflow and velocity. When the force generated by the oropharyngeal dilator and abductor muscles is exceeded by a critical amount, the negative airway pressure increases and facilitates compliant pharyngeal tissues to collapse by acting as a Starling resistor.⁷⁰ This mechanism is recognized in disturbed sleep caused by nasal obstruction.⁷¹

Furthermore, several inflammatory mediators were reported to be associated with poor sleep, including histamine, CysLTs, IL-1 β , IL-4, IL-10, bradykinin, and substance P.²⁷ Increased mediators induced by rhinitis may contribute to disturbances of sleep architecture in OSA patients.²⁵ Histamine, a relevant modulator of the sleep-wake cycle, also influences mechanisms related to arousal, cognition, and memory.²⁵ The typical increase in histaminergic function associated with different allergic conditions and the evidence of sleep architecture improvements with anti-histaminic therapy reinforce the significance of the relationship between AR and sleep.⁷² An exploratory study demonstrated that levels of IL-1 β , IL-4, and IL-10 were higher in AR patients, compared with non-allergic patients, and correlated with increased latency to REM sleep, decreased time spent in REM sleep and decreased latency to sleep onset.⁷³ Likewise, OSA has also been shown to be associated with increased cytokines such as IL-1, IL-4, IL-6, and Tumour Necrosis Factors (TNF) that promote T helper type 2 (Th2) cell phenotypes and enhance inflammation leading to nasal congestion.⁷⁴⁻⁷⁶ One additional consequence of the intermittent hypoxia and sleep fragmentation is oxidative stress, with reactive oxygen species production and lipid peroxidation being found to increase in OSA patients.⁷⁴ The association between chronic inflammation and oxidative stress is well documented. Oxidative stress results from the imbalance between the oxidative forces and the antioxidant defense systems, which is believed to favor an oxidative injury that has been implicated in the pathogenesis of AR.⁷⁷

Th17/Treg ratios are elevated in OSA patients and are positively associated with the Apnea-Hypopnea Index (AHI).²⁵ Similarly, it was shown that the Th17/Treg ratio was significantly higher in OSA children with AR compared to OSA children with ATH but without AR.⁷⁸ Th17 positive cells play critical roles in the development of autoimmunity and allergic reactions through IL-17 production. Treg cells orchestrate the overall immune response and play a role in immune tolerance. Th17 and Treg imbalance caused by AR and OSA,^{79,80} respectively, may mutually promote each other and lead to further imbalance. Thus, regardless of which pathological changes appear first, both AR and OSA will facilitate manifestations of Th17/Treg imbalance.⁷⁸

The trigeminocardiac reflex (TCR) has been advanced as a putative mechanism linking AR to sleep disorders, as TCR can be activated by stimulation of the trigeminal nerve anywhere along its course, eliciting from mild to severe negative cardiorespiratory changes, such as bradycardia, asystole, hypotension or apnea, either due to sympathetic withdrawal or to parasympathetic overactivity via the vagus nerve.⁸¹ Further, AR is known to induce neuronal hyperresponsiveness of the upper airways to stimuli that activate nasal afferents.

Dysregulation of nasal neural pathways plays a relevant role in the pathogenesis of rhinitis and causes nasal, respiratory, and cardiac changes. Nasal discharge or nasal congestion can be potential stimuli for nasal trigeminal afferents in rhinitis and TCR, and may also contribute as a mechanism underlying the pathogenesis of sleep disorders in AR.⁸¹

When examining the impact of AR on all dimensions of sleep quality, it is important to consider the potential impact of co-existing factors such as anxiety and depression or of frequent comorbidities such as asthma. Actually, in early studies, it was estimated that 15 to 40% of AR patients had asthma, and 6% to 85% of asthmatic patients reported AR.^{82,83}

The authors would be remiss if we did not mention that sleep quality exerts an important influence on host immune system homeostasis. A night of sleep deprivation or 5 days of sleep restriction will alter the TH1/TH2 balance.⁸⁴ Furthermore, environmental (e.g., allergens and irritants,⁸⁵ individual (e.g., sleep hygiene and adherence to treatment) and family/cultural factors (e.g., race, ethnicity, factors related to urban poverty, beliefs and family sleep practices), in combination or on their own, exert influence on both allergic diseases and sleep and need to be considered when working with patients who have allergic disorders.⁸⁶

Treatment of AR and effects on sleep

Although sleep is crucial for optimal child growth, development, and learning ability, a survey of Pediatric Allergies in America indicated that 40% of parents claimed that nasal allergy affected their child sleep: 32% claimed their child had difficulty falling asleep, 26% claimed their child awakened at night due to sleep problems, and 29% claimed their children had a lack of good sleep.⁶ This relationship between allergic diseases and sleep disorders in the pediatric age has also been confirmed in some Latin American countries.⁸⁷

Therapies that reduce nasal congestion may improve sleep and daytime sleepiness and, thereby, improve quality of life. In addition, those therapies that reduce inflammation, often those that also decrease congestion, may have a positive influence by reducing the levels of inflammatory mediators, such as TNF, which can cause constitutional symptoms. However, not all treatments for AR are effective in this respect.⁸⁸ The authors also point out that data regarding the effect of allergen avoidance on sleep in AR patients is limited.⁸⁹ The effects of medication and isotonic or hypertonic saline were evaluated to assess how AR treatment may affect sleep and SDB.^{75,89,90} These trials included intranasal corticosteroids, leukotriene receptor antagonists, and topical nasal decongestants. Noteworthy, there were few studies in children, and relevant evidence is limited since most of the studies assessing treatments effects included a small number of subjects and were not conducted in a randomized double-blinded placebo-controlled fashion.⁷⁵

Topical decongestants reduce nasal congestion but may have adverse effects on sleep because of their stimulatory effects and also because of the risk of rhinitis medicamentosa ("rebound" congestion), such that they should not be used for extended periods of time.⁸⁸ Sedating antihistamines are generally contraindicated, particularly for those children already experiencing daytime somnolence, fatigue, and functional impairments. Non-sedating oral antihistamines are widely used to treat AR and relieve nasal symptoms such as rhinorrhea, sneezing, and pruritus but have less effect on nasal congestion.⁸⁸ Some studies have shown an improvement in sleep and quality of life following therapy with an oral or topical antihistamine in adults.⁸⁸ Data regarding intranasal antihistamines and effects on sleep are mixed, with some studies demonstrating improvement in quality of sleep and proportion of mouth breathing, while others indicate increased daytime somnolence as a side effect of these agents.⁹¹

A prospective randomized, double-blind controlled trial of PSG-diagnosed OSA in 57 children, ages 2-10 years, with or without rhinitis, who were treated with either oral montelukast or placebo, demonstrated that montelukast for 16 weeks effectively reduced the severity of OSA.⁹² Montelukast markedly improved quality of life in children with mild OSA.⁹³ Nevertheless, the current concerns about potential undesirable side-effects with this drug raise substantial barriers to initiation of such therapy in the age group.⁹⁴ In recent years, there has been an increased focus on neuropsychiatric adverse drug events (ADE) associated with leukotriene antagonists, owing to a 2008 US FDA alert on possible neuropsychiatric ADE.⁹⁵ A 2016 review of pediatric psychiatric disorders associated with Montelukast using the VigiBase demonstrated age-variant neuropsychiatric ADE, with infants and children developing sleep disturbances and predisposing to depression/anxiety and psychotic reactions in adolescents.⁹⁶

Multiple placebo-controlled clinical trials have shown that intranasal corticosteroids are effective against the most common symptoms of AR, including sneezing, itching, rhinorrhea, and congestion in both adults and children.⁸⁹ Intranasal corticosteroids are considered first-line therapy when nasal congestion is a major symptom. Small studies in children with AR found that intranasal budesonide reduces nasal congestion, subjective daytime somnolence, and fatigue and improves patient sleep and life quality.⁸⁸ Indeed, during treatment of children with AR with intranasal budesonide, nasal symptoms decreased, objective sleep studies were improved, and subjective symptoms relating to poor sleep quality were diminished.⁹⁷ Treatment with intranasal budesonide may also improve concentration and performance at school in children with rhinitis.⁹⁸

Besides the effects of topical corticosteroids on rhinitis control, ATH and OSA may respond favorably to anti-inflammatory agents such as corticosteroids and montelukast via their lymphocytic, anti-proliferative or anti-inflammatory effects, all of which might reduce adenotonsillar hypertrophy. Intranasal corticosteroids reduce cellular proliferation and the production of pro-inflammatory cytokines in a tonsil and adenoid mixed-cell culture system.⁹⁹ As a corollary of such findings, randomized controlled trials of intranasal corticosteroids have resulted in significant improvements or resolution of mild OSA in children.^{100,101}

Adequate treatment of AR could prevent the occurrence of OSA, reduce the severity of existing OSA, and potentially prevent the emergence of the maxillofacial growth abnormalities that predispose children to OSA.¹⁰² Multivariate analysis has also identified rhinitis as an independent predictor for the failure of AT and the presence of residual OSA.¹⁰³

AR treatment drugs such as Azelastine nasal spray and Mometasone Furoate Aqueous Nasal Spray were found to be useful in decreasing adenoid pad size and the severity of symptoms related to adenoidal hypertrophy in children.^{104,105} So far, therapeutic studies with azelastine hydrochloride and fluticasone propionate in a single delivery system using objective sleep metrics in children and adolescents with AR are still lacking. Therefore, the advantage of the combination treatment, although promising, remains uncertain.

Concerns and future directions

Sleep problems are a commonly associated morbidity in people with AR, and disrupted sleep homeostasis has many well-documented consequences; patients with AR have increased daytime and night-time symptoms related to sleep disorders; although the relative strength of studies results is low, AR has a significant association with both snoring and OSA among children, and intranasal corticosteroids or montelukast improve both AR and SDB.

Although a substantial number of studies have analyzed the relationship between AR and OSA either in the context of pathophysiological, epidemiological or therapeutic pathways, several methodological issues turn difficult the generalization of some important findings. In this context, one important issue is that most of the available literature support findings in adults which is therefore hard to transpose to pediatric ages. Also, the use of distinct assessment tools leads to a harder interpretation and uniformization of findings. Future research would benefit from increased attention on those topics.

Further studies are needed to more compellingly demonstrate or refute the impact of AR on sleep and vice-versa with improved delineation of the criteria for AR definition and the use of PSG for sleep assessments; to substantiate that allergic sensitization might increase SDB severity in children and verify the effects of AR treatment in children with sleep disorders, using large double-blind placebo-controlled clinical trials.

Conclusions

Although conflictive, there is currently a great body of evidence precluding firm statements on the interactive roles between AR and sleep. Still, identifying and treating rhinitis in clinical practice have been pointed out as an important step toward improving the symptoms and preventing sleep quality deterioration in children with SDB/OSA before and after undergoing AT. The future of pediatric clinical research will certainly benefit from further exploration of such a relationship between AR and sleep disorders which seems to persist as a clinically relevant interaction in modern days.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Abdelhafeez M. Effectiveness of intranasal steroids on rhinitis symptoms, sleep quality, and quality of life in patients with perennial allergic rhinitis. *Eur Arch Otorhinolaryngol*. 2021 Apr 17. <https://doi.org/10.1007/s00405-021-06818-1>. Epub ahead of print.
- Gelfand EW. Inflammatory mediators in allergic rhinitis. *J Allergy Clin Immunol*. 2004;114:S135–8.
- Li Z, Celestin J, Lockey RF. Pediatric sleep apnea syndrome: an update. *J Allergy Clin Immunol Pract*. 2016;4:852–61.
- Liu J, Zhang X, Zhao Y, Wang Y. The association between allergic rhinitis and sleep: a systematic review and meta-analysis of observational studies. *PLoS One*. 2020;15:e0228533.
- Öçal R, Bayar Muluk N, MULLOL J. Epidemiology of allergic rhinitis. In: Cingi C, Bayar Muluk N, eds. *All around the nose*, Cham: Springer; 2020:297–301.
- Meltzer EO, Blaiss MS, Derebery MJ, Mahr TA, Gordon BR, Sheth KK, et al. Burden of allergic rhinitis: results from the pediatric allergies in america survey. *J Allergy Clin Immunol*. 2009;124:543–70.
- Mallol J, Crane J, von Mutius E, Odhiambo J, Keil U, Stewart A, et al. The international study of asthma and allergies in childhood (ISAAC) phase three: a global synthesis. *Allergol Immunopathol (Madr)*. 2013;41:73–85.
- Léger D, Annesi-Maesano I, Carat F, Rugina M, Chanal I, Pribil C, et al. Allergic rhinitis and its consequences on quality of sleep: an unexplored area. *Arch Intern Med*. 2006;166:1744–8.
- Asher MI, Montefort S, Björkstén B, Lai CK, Strachan DP, Weiland SK, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC phases one and three repeat multicountry cross-sectional surveys [published correction appears in *Lancet* 2007 Sep 29;370(9593):1128]. *Lancet*. 2006;368:733–43.
- Galli SJ, Tsai M. IgE and mast cells in allergic disease. *Nat Med*. 2012;18:693–704.
- Gauvreau GM, El-Gammal AI, O'Byrne PM. Allergen-induced airway responses. *Eur Respir J*. 2015;46:819–31.
- Rosenwasser LJ. Current understanding of the pathophysiology of allergic rhinitis. *Immunol Allergy Clin North Am*. 2011;31:433–9.
- Nakao A, Nakamura Y, Shibata S. The circadian clock functions as a potent regulator of allergic reaction. *Allergy*. 2015;70:467–73.
- Skoner DP. Allergic rhinitis: definition, epidemiology, pathophysiology, detection, and diagnosis. *J Allergy Clin Immunol*. 2001;108:S2–8.
- Reinberg A, Gervais P, Levi F, Smolensky M, Del Cerro L, Ugolini C. Circadian and circannual rhythms of allergic rhinitis: an epidemiologic study involving chronobiologic methods. *J Allergy Clin Immunol*. 1988;81:51–62.
- Wang X, Reece SP, Van Scott MR, Brown JM. A circadian clock in murine bone marrow-derived mast cells modulates IgE-dependent activation in vitro. *Brain Behav Immun*. 2011;25:127–34.
- Dickmeis T. Glucocorticoids and the circadian clock. *J Endocrinol*. 2008;200:3–22.
- Cheng FL, An YF, Han ZQ, Li C, Li ZQ, Yang PC, et al. Period2 gene regulates diurnal changes of nasal symptoms in an allergic rhinitis mouse model. *Int Forum Allergy Rhinol*. 2020;10:1236–48.
- Nakao A. Circadian regulation of the biology of allergic disease: clock disruption can promote allergy. *Front Immunol*. 2020;11:1237.
- Smolensky MH, Lemmer B, Reinberg AE. Chronobiology and chronotherapy of allergic rhinitis and bronchial asthma. *Adv Drug Deliv Rev*. 2007;59:852–82.
- Aoyagi M, Watanabe H, Sekine K, Nishimuta T, Konno A, Shimajo N, et al. Circadian variation in nasal reactivity in children with allergic rhinitis: correlation with the activity of eosinophils and basophilic cells. *Int Arch Allergy Immunol*. 1999;120:595–9.
- Berson SR, Klimczak JA, Prezio EA, Abraham MT. House dust mite related allergic rhinitis and REM sleep disturbances. *Am J Otolaryngol*. 2020;41:102709.
- Stormark KM, Fosse HE, Pallesen S, Hysing M. The association between sleep problems and academic performance in primary school-aged children: findings from a Norwegian longitudinal population-based study. *PLoS One*. 2019;14:e0224139.

24. Lumeng JC, Chervin RD. Epidemiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc.* 2008;5:242–52.
25. Zheng M, Wang X, Zhang L. Association between allergic and non-allergic rhinitis and obstructive sleep apnea. *Curr Opin Allergy Clin Immunol.* 2018;18:16–25.
26. Banasiak NC. Understanding the relationship between asthma and sleep in the pediatric population. *J Pediatr Health Care.* 2016;30:546–50.
27. Ferguson BJ. Influences of allergic rhinitis on sleep. *Otolaryngol Head Neck Surg.* 2004;130:617–29.
28. Lin SY, Melvin TA, Boss EF, Ishman SL. The association between allergic rhinitis and sleep-disordered breathing in children: a systematic review. *Int Forum Allergy Rhinol.* 2013;3:504–9.
29. Chng SY, Goh DY, Wang XS, Tan TN, Ong NB. Snoring and atopic disease: a strong association. *Pediatr Pulmonol.* 2004;38:210–6.
30. Tafur A, Chérrez-Ojeda I, Patiño C, Gozal D, Rand C, Ronnie M, et al. Rhinitis symptoms and habitual snoring in Ecuadorian children. *Sleep Med.* 2009;10:1035–9.
31. Anuntaseree W, Rookkapan K, Kuasirikul S, Thongsuksai P. Snoring and obstructive sleep apnea in Thai school-age children: prevalence and predisposing factors. *Pediatr Pulmonol.* 2001;32:222–7.
32. Ng DK, Chan CH, Hwang GY, Chow PY, Kwok KL. A review of the roles of allergic rhinitis in childhood obstructive sleep apnea syndrome. *Allergy Asthma Proc.* 2006;27:240–2.
33. Lofaso F, Coste A, d'Ortho MP, Zerah-Lancner F, Delclaux C, Goldenberg F, et al. Nasal obstruction as a risk factor for sleep apnoea syndrome. *Eur Respir J.* 2000;16:639–43.
34. Giraldo-Cadavid LF, Perdomo-Sanchez K, Córdoba-Gravini JL, Escamilla MI, Suarez M, Gelvez N, Gozal D, et al. Allergic rhinitis and osa in children residing at a high altitude. *Chest.* 2020;157:384–93.
35. Huseni S, Gutierrez MJ, Rodriguez-Martinez CE, Nino CL, Perez GF, Panchar K, et al. The link between rhinitis and rapid-eye-movement sleep breathing disturbances in children with obstructive sleep apnea. *Am J Rhinol Allergy.* 2014;28:56–61.
36. Tamanyan K, Walter LM, Davey MJ, Nixon GM, Horne RS, Biggs SN. Risk factors for obstructive sleep apnoea in Australian children. *J Paediatr Child Health.* 2016;52:512–7.
37. Liu J, Wu Y, Wu P, Xu Z, Ni X. Analysis of the impact of allergic rhinitis on the children with sleep disordered breathing. *Int J Pediatr Otorhinolaryngol.* 2020;138:110380.
38. Di Francesco RC, Alvarez J. Allergic rhinitis affects the duration of rapid eye movement sleep in children with sleep-disordered breathing without sleep apnea. *Int Forum Allergy Rhinol.* 2016;6:465–71.
39. Nosetti L, Piacentini G, Macchi A, De Bernardi F, Simoncini D, Nicoloso M, et al. Nasal cytology in children with primary snoring and obstructive sleep apnoea syndrome. *Int J Pediatr Otorhinolaryngol.* 2019;122:133–7.
40. Xu Z, Wu Y, Tai J, Feng G, Ge W, Zheng L, et al. Risk factors of obstructive sleep apnea syndrome in children. *J Otolaryngol Head Neck Surg.* 2020;49:11.
41. Weinstock TG, Rosen CL, Marcus CL, Garetz S, Mitchell RB, Amin R, et al. Predictors of obstructive sleep apnea severity in adenotonsillectomy candidates. *Sleep.* 2014;37:261–9.
42. Kalra M, Lemasters G, Bernstein D, Wilson K, Levin L, Cohen A, et al. Atopy as a risk factor for habitual snoring at age 1 year. *Chest.* 2006;129:942–6.
43. Zhang J, Zhao J, Chen M, Liu S, Zhang X, Zhang F, et al. Airway resistance and allergic sensitization in children with obstructive sleep apnea hypopnea syndrome. *Pediatr Pulmonol.* 2016;51:426–30.
44. Kramer MF, De La Chaux R, Dreher A, Pfrogner E, Rasp G. Allergic rhinitis does not constitute a risk factor for obstructive sleep apnea syndrome. *Acta Otolaryngol.* 2001;121:494–9.
45. Shen L, Lin Z, Lin X, Yang Z. Risk factors associated with obstructive sleep apnea-hypopnea syndrome in Chinese children: a single center retrospective case-control study. *PLoS One.* 2018;13:e0203695.
46. Loekmanwidjaja J, Carneiro ACF, Nishinaka MLT, Munhoes DA, Benezoli G, Wandalsen GF, et al. Sleep disorders in children with moderate to severe persistent allergic rhinitis. *Braz J Otorhinolaryngol.* 2018;84:178–84.
47. Lai PH, Yang PS, Lai WY, Lin CL, Hsu CY, Wei CC. Allergic rhinitis and the associated risk of nocturnal enuresis in children: a population-based cohort study. *Int Forum Allergy Rhinol.* 2018;8:1260–6.
48. Bilgiliyoz Filiz M, Filiz S, Baran RT, Çakır T, Koldaş Doğan Ş, Parlak M, et al. Restless legs syndrome in children with allergic rhinitis: a comparative study on frequency, severity and sleep quality. *Turk J Phys Med Rehabil.* 2018;64:198–204.
49. Tsai JD, Chen HJ, Ku MS, Chen SM, Hsu CC, Tung MC, et al. Association between allergic disease, sleep-disordered breathing, and childhood nocturnal enuresis: a population-based case-control study. *Pediatr Nephrol.* 2017;32:2293–301.
50. Nguyen-Hoang Y, Nguyen-Thi-Dieu T, Duong-Quy S. Study of the clinical and functional characteristics of asthmatic children with obstructive sleep apnea. *J Asthma Allergy.* 2017;10:285–92.
51. Poachanukoon O, Kitcharoensakkul M. Snoring and sleep problems in children with and without allergic rhinitis: a case control study. *J Med Assoc Thai.* 2015;98:S138–44.
52. Ng DK, Chan CH, Ng EP. Natural history of snoring in Hong Kong adolescents. *J Paediatr Child Health.* 2014;50:596–604.
53. Fadzil Abdullah AA, Jamalludin AR, Norrashidah AW, Norzila MZ, Asiah Kassim K, Rus Anida A, et al. Prevalence of sleep disordered breathing symptoms among Malay school children in a primary school in Malaysia. *Med J Malaysia.* 2012;67:181–5.
54. So HK, Li AM, Au CT, Zhang J, Lau J, Fok TF, et al. Night sweats in children: prevalence and associated factors. *Arch Dis Child.* 2012;97:470–3.
55. Ishman SL, Smith DF, Benke JR, Nguyen MT, Lin SY. The prevalence of sleepiness and the risk of sleep-disordered breathing in children with positive allergy test. *Int Forum Allergy Rhinol.* 2012;2:139–43.
56. Bhattacharjee R, Kheirandish-Gozal L, Spruyt K, Mitchell RB, Promchiarak J, Simakajornboon N, et al. Adenotonsillectomy outcomes in treatment of obstructive sleep apnea in children: a multicenter retrospective study. *Am J Respir Crit Care Med.* 2010;182:676–83.
57. Urschitz MS, Brockmann PE, Schlaud M, Poets CF. Population prevalence of obstructive sleep apnoea in a community of German third graders. *Eur Respir J.* 2010;36:556–68.
58. Li S, Jin X, Yan C, Wu S, Jiang F, Shen X. Habitual snoring in school-aged children: environmental and biological predictors. *Respir Res.* 2010;11:144.
59. Sogut A, Yilmaz O, Dinc G, Yuksel H. Prevalence of habitual snoring and symptoms of sleep-disordered breathing in adolescents. *Int J Pediatr Otorhinolaryngol.* 2009;73:1769–73.
60. Liukkonen K, Virkkula P, Aronen ET, Kirjavainen T, Pitkäranta A. All snoring is not adenoids in young children. *Int J Pediatr Otorhinolaryngol.* 2008;72:879–84.
61. Petry C, Pereira MU, Pitrez PM, Jones MH, Stein RT. The prevalence of symptoms of sleep-disordered breathing in Brazilian schoolchildren. *J Pediatr (Rio J).* 2008;84:123–9.
62. Sogut A, Altin R, Uzun L, Ugur MB, Tomac N, Acun C, et al. Prevalence of obstructive sleep apnea syndrome and associated symptoms in 3–11-year-old Turkish children. *Pediatr Pulmonol.* 2005;39:251–6.
63. Ng DK, Kwok KL, Cheung JM, Leung SY, Chow PY, Wong WH, et al. Prevalence of sleep problems in Hong Kong primary school children: a community-based telephone survey. *Chest.* 2005;128:1315–23.

64. Ibáñez V, Silva J, Cauli O. A survey on sleep questionnaires and diaries. *Sleep Med.* 2018;42:90–6.
65. Silva FG, Silva CR, Braga LB, Neto AS. Portuguese Children's Sleep Habits Questionnaire - validation and cross-cultural comparison. *J Pediatr (Rio J).* 2014;90:78–84.
66. Kaditis AG, Finder J, Alexopoulos EI, Starantzis K, Tanou K, Gampeta S, et al. Sleep-disordered breathing in 3,680 Greek children. *Pediatr Pulmonol.* 2004;37:499–509.
67. Evcimik MF, Dogru M, Cirik AA, Nepesov MI. Adenoid hypertrophy in children with allergic disease and influential factors. *Int J Pediatr Otorhinolaryngol.* 2015;79:694–7.
68. Huo Z, Shi J, Shu Y, Xiang M, Lu J, Wu H. The relationship between allergic status and adenotonsillar regrowth: a retrospective research on children after adenotonsillectomy. *Sci Rep.* 2017;7:46615.
69. Mukundan TH. Screening for allergic disease in a child with sleep disorder and screening for sleep disorder in allergic disease. In: Fishbein A, Sheldon SH, eds. *Allergy and sleep*, Cham: Springer International Publishing; 2019:77–85.
70. Marcus CL, McColley SA, Carroll JL, Loughlin GM, Smith PL, Schwartz AR. Upper airway collapsibility in children with obstructive sleep apnea syndrome. *J Appl Physiol.* 1994;77:918–24. 1985.
71. Kakumanu S, Glass C, Craig T. Poor sleep and daytime somnolence in allergic rhinitis. *Am J Respir Crit Care Med.* 2002;1:195–200.
72. Thakkar MM. Histamine in the regulation of wakefulness. *Sleep Med Rev.* 2011;15:65–74.
73. Krouse HJ, Davis JE, Krouse JH. Immune mediators in allergic rhinitis and sleep. *Otolaryngol Head Neck Surg.* 2002;126:607–13.
74. Maniaci A, Iannella G, Cocuzza S, Vicini C, Magliulo G, Ferlito S, et al. Oxidative stress and inflammation biomarker expression in obstructive sleep apnea patients. *J Clin Med.* 2021;10:277.
75. Chirakalwasan N, Ruxrungtham K. The linkage of allergic rhinitis and obstructive sleep apnea. *Asian Pac J Allergy Immunol.* 2014;32:276–86.
76. Kheirandish-Gozal L, Gozal D. Obstructive sleep apnea and inflammation: proof of concept based on two illustrative cytokines. *Int J Mol Sci.* 2019;20:459.
77. Shiefa S, Ashalatha R, Anjali R. Increased oxidative stress and altered antioxidants status in patients with chronic allergic rhinitis. *Adv Biosci Biotechnol.* 2012;03:951–6.
78. Ni K, Zhao L, Wu J, Chen W, HongyaYang Li X. Th17/Treg balance in children with obstructive sleep apnea syndrome and the relationship with allergic rhinitis. *Int J Pediatr Otorhinolaryngol.* 2015;79:1448–54.
79. Tan HL, Gozal D, Wang Y, Bandla HP, Bhattacharjee R, Kulkarni R, et al. Alterations in circulating T-cell lymphocyte populations in children with obstructive sleep apnea. *Sleep.* 2013;36:913–22.
80. Tan HL, Gozal D, Samiei A, Bhattacharjee R, Wang Y, Ramirez HM, et al. T regulatory lymphocytes and endothelial function in pediatric obstructive sleep apnea. *PLoS One.* 2013;8:e69710.
81. Bindu B, Singh GP, Chowdhury T, Schaller B. Rhinitis and sleep disorders: the trigeminocardiac reflex link? *Med Hypotheses.* 2017;103:96–9.
82. Kim H, Bouchard J, Renzix P. The link between allergic rhinitis and asthma: a role for antileukotrienes? *Can Respir J.* 2008;15:91–8.
83. Brożek JL, Bousquet J, Agache I, Agarwal A, Bachert C, Bosnic-Anticevich S, et al. Allergic rhinitis and its impact on asthma (ARIA) guidelines-2016 revision. *J Allergy Clin Immunol.* 2017;140:950–8.
84. Nunes JO, Apostolico JS, Andrade DAG, Ruiz FS, Fernandes ER, Andersen ML, et al. Sleep deprivation predisposes allergic mice to neutrophilic lung inflammation. *J Allergy Clin Immunol.* 2018;141:1018–27. e4.
85. Naclerio R, Ansotegui IJ, Bousquet J, Walter Canonica G, D'Amato G, Rosario N, et al. International expert consensus on the management of allergic rhinitis (AR) aggravated by air pollutants: impact of air pollution on patients with AR: current knowledge and future strategies. *World Allergy Organ J.* 2020;13:100106.
86. Koinis-Mitchell D, Craig T, Esteban CA, Klein RB. Sleep and allergic disease: a summary of the literature and future directions for research. *J Allergy Clin Immunol.* 2012;130:1275–81.
87. Urrutia-Pereira M, Solé D, Chong Neto HJ, Acosta V, Cepeda AM, Álvarez-Castelló M, et al. Sleep disorders in Latin-American children with asthma and/or allergic rhinitis and normal controls. *Allergol Immunopathol (Madr).* 2017;45:145–51.
88. Lunn M, Craig T. Rhinitis and sleep. *Sleep Med Rev.* 2011;15:293–9.
89. Correia C, Baroody FM. Allergic rhinitis and sleep: approaches to management. In: Fishbein A, Sheldon SH, eds. *Allergy and sleep*, Cham: Springer International Publishing; 2019:271–92.
90. Marcuccio G, Di Bari M, Precenzano F, Operto FF, Bitetti I, Motta G, et al. Relationship between sleep quality and rhinitis in children: role of medical treatment with isotonic and hypertonic salines. *Minerva Pediatr (Torino).* 2021;73:301–6.
91. McLean HA, Urton AM, Driver HS, Tan AK, Day AG, Munt PW, et al. Effect of treating severe nasal obstruction on the severity of obstructive sleep apnoea. *Eur Respir J.* 2005;25:521.
92. Kheirandish-Gozal L, Bandla HP, Gozal D. Montelukast for children with obstructive sleep apnea: results of a double-blind, randomized, placebo-controlled trial. *Ann Am Thorac Soc.* 2016;13:1736–41.
93. Bluher AE, Brawley CC, Cunningham TD, Baldassari CM. Impact of montelukast and fluticasone on quality of life in mild pediatric sleep apnea. *Int J Pediatr Otorhinolaryngol.* 2019 Oct;125:66–70.
94. Kovesi T. Neuropsychiatric side effects of montelukast. *J Pediatr.* 2019;212:48.
95. Leung JS, Johnson DW, Sperou AJ, Crotts J, Saude E, Hartling L, et al. A systematic review of adverse drug events associated with administration of common asthma medications in children. *PLoS One.* 2017;12:e0182738.
96. Aldea Perona A, Garcia-Saiz M, Sanz Alvarez EA-Ohoo. Psychiatric disorders and montelukast in children: a disproportionality analysis of the VigiBase®. *Drug Safety.* 2016;39:69–78.
97. Mansfield LE, Diaz G, Posey CR, Flores-Neder J. Sleep disordered breathing and daytime quality of life in children with allergic rhinitis during treatment with intranasal budesonide. *Ann Allergy Asthma Immunol.* 2004;92:240–4.
98. Kamenov S, Kamenov B, Moskovljevic J, Zivkovic M, Zaharov T. Intranasal budesonide treatment of rhinosinusitis improves quality of life in school children. In: Presented at: World Allergy Congress, Munich, Germany; 2005. June 26eJuly 1.
99. Sakarya EU, Bayar Muluk N, Sakalar EG, Senturk M, Aricigil M, Bafaqeeh SA, et al. Use of intranasal corticosteroids in adenotonsillar hypertrophy. *J Laryngol Otol.* 2017;131:384–90.
100. Gozal D, Tan HL, Kheirandish-Gozal L. Treatment of obstructive sleep apnea in children: handling the unknown with precision. *J Clin Med.* 2020;9:888.
101. Kheirandish-Gozal L, Bhattacharjee R, Bandla HPR, Gozal D. Antiinflammatory therapy outcomes for mild OSA in children. *Chest.* 2014;146:88–95.
102. Smith RM, Gonzalez C. The relationship between nasal obstruction and craniofacial growth. *Pediatr Clin North Am.* 1989;36:1423–34.
103. Alsufyani N, Isaac A, Witmans M, Major P, El-Hakim H. Predictors of failure of DISE- directed adenotonsillectomy in children

- with sleep disordered breathing. *J Otolaryngol Head Neck Surg.* 2017;46:37.
104. Berkiten G, Kumral TL, Çakır O, Yıldırım G, Salturk Z, Uyar Y, et al. Effectiveness of azelastine nasal spray in the treatment of adenoidal hypertrophy in children. *Hippokratia.* 2014;18: 340–5.
105. Berlucchi M, Salsi D, Valetti L, Parrinello G, Nicolai P. The role of mometasone furoate aqueous nasal spray in the treatment of adenoidal hypertrophy in the pediatric age group: preliminary results of a prospective, randomized study. *Pediatrics.* 2007;119: e1392–7.