



ORIGINAL ARTICLE

Parental, gestational, and early-life exposure to indoor environmental hazardous factors on allergic rhinitis among preschool children in Urumqi City: a case-control study



Tingting Wang ^{a,1}, Haonan Shi ^{a,1}, Huizhen Qi ^{b,1}, Lan Jiang ^{c,1}, Yuting Lin ^{c,1}, Jian Yao ^d, Junwen Lu ^d, Qi Yan ^{d,*}, Aiyong Zhu ^{a,*}, Xuan Yao ^{e,*}

^a Shanghai University of Medicine & Health Sciences, School of Nursing & Health Management, Shanghai, China

^b The Fifth Affiliated Hospital of Xinjiang Medical University, Department of Neurology, Urumqi, China

^c Xinjiang Uyghur Autonomous Region Maternal and Child Health Hospital, Department of Laboratory Medicine, Urumqi, China

^d Xinjiang Medical University, School of Public Health, Urumqi, China

^e Shanghai University, College of Medicine, Shanghai, China

Received 20 June 2022; accepted 15 September 2022

Available online 27 January 2023

KEYWORDS

Allergic rhinitis;
Preschool children;
Indoor environment

Abstract

Objective: Allergic rhinitis (AR) is a disease associated with impaired quality of life and heredity. This study aimed to investigate the association of allergic rhinitis in preschool children with exposure to indoor environment-related factors early in life.

Methods: In August 2019, the authors implemented a study among 2020 preschool children in Urumqi City using a case-control design. The study included parental reports for the occurrence of AR in children, parental history of respiratory disease, and indoor environmental correlates of maternal exposure from 1 year prior to pregnancy until the child's age of 0-1 year.

Results: Mode of birth (cesarean section) (OR = 1.31, 95% CI = 1.02~1.67), father with AR (OR = 2.67, 95% CI = 2.08~3.44), mother with AR (OR = 3.70, 95% CI = 2.88~4.74), mother with asthma (OR = 3.11, 95% CI = 1.18~8.20), and mother with newly purchased furniture in the parents' residence during pregnancy (OR = 1.49, 95% CI = 1.03~2.14) were risk factors for AR in children.

Conclusions: The focus of allergic rhinitis should be on children with a family history of AR and asthma and cesarean delivery. Primary prevention efforts for AR in preschool children are avoiding exposure of children to indoor environmental hazardous factors early in life.

© 2023 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mails: 13579939479@163.com (Q. Yan), zhuay@sumhs.edu.cn (A. Zhu), 389903786@qq.com (X. Yao).

¹ These authors contributed equally to this work.

Introduction

Allergic rhinitis (AR) refers to a noninfectious inflammatory disease of the nasal mucosa involving multiple immunocompetent cells and cytokines mediated by IgE after allergen exposure in atopic individuals.¹ Most people ignore the harms posed by AR because of its milder symptoms, but several studies have confirmed that AR causes a range of serious harms. An important study from the Chinese lung health study group published in *The Lancet* in 2019² showed that AR is an important risk factor for asthma in adults. In addition, symptoms of AR adversely affect patients' work and quality of life, and in severe cases, sleep-disordered breathing can occur.³ AR also causes symptoms such as reduced learning ability and poor concentration in children and adolescents.⁴ A 2012 study from 10 cities in China involving an indoor environment and child health (CCHH) showed a prevalence of physician-diagnosed AR ranging from 2.2% to 23.9% in Chinese preschool children, including 9.8% in Urumqi City.⁵ Since Urumqi City is a multiethnic aggregation site, there are large differences in lifestyle habits among ethnic groups, and the climate is dry, which may affect the transmission of allergens within the environment. It is, therefore, necessary to carry out a large epidemiological survey study of AR in preschool children as soon as possible. A 2019 study suggested that exposure to environmental risk factors early in life may increase the incidence of AR in children.⁶

For the reasons set out above, the prevalence of AR among preschool-aged children and mothers' residential environmental factors from preconception to the age of 1 year in 60 kindergartens in Xinshi District, Shayibake District, Tianshan District, Shuimogou District, Toutunhe District, and Midong District were investigated in August 2019. Exploring the association between exposure to indoor environment-related factors and AR in children during this period provides a scientific and theoretical basis for the prevention of AR in children.

Methods

Investigated subjects

In August 2019, the authors used stratified random sampling, six administrative districts were first selected in the Urumqi, which were Xinshi District, Shayibake District, Tianshan District, Shuimogou District, Toutunhe District, and Midong District. Then 8-12 kindergartens were randomly selected in each district. A total of 60 kindergartens were selected, and children with AR who diagnosed by doctors were investigated. In the same period, children who were sex -, age -, and ethnicity matched and had not been diagnosed with AR in the same kindergartens were selected as controls in a 1:4 ratio of the case: control. Every control child was included in this study with equal probability. This study was approved by the ethics committee of the Hospital, and the parents of the investigated children provided written informed consent.

Investigation contents and methods

Questionnaires were developed for all children in the case group and control group; the questionnaire contents were

adapted from the questionnaires used in the CCHH survey study and were slightly revised according to the specific context of Urumqi City.

The specific survey included:

- (1) Demographic characteristics, including sex, ethnicity, and age at the home address.
- (2) Feeding status, including whether they were only children, whether they were breastfed, and duration of breastfeeding.
- (3) Prevalence of AR in children and family members, including AR in children and parents and AR-related symptoms.
- (4) Dwelling environment, including housing type, renovation situation, and furniture purchase situation.
- (5) Living habits, including indoor pets and house plants, smoking, air conditioning, and the use of a new air filtering system.
- (6) Dietary habits, including meal types and times.

Before the survey began, the subject group first had contact with the Education Bureau of Urumqi City and the kindergarten teachers, and uniform professional training was provided to the responsible teachers involved in the survey class before the survey. Consent for the questionnaire to be administered to the parents of the children was given in their kindergartens by the preschool teachers, and the questionnaires were completed by the parents or other guardians of the children at home. They were asked to complete the form within 1 week and submit it to the responsible teacher at the kindergarten, who then returned the questionnaire and uniformly sent it to the Education Bureau of Urumqi City. All questionnaires were reviewed by more than 2 trained subject team members, and ineligible questionnaires were excluded.

Determination of outcomes

- (1) AR: parent reports that the child has had at least 1 occasion in which AR has been definitively diagnosed by a physician, with the age of onset determined by the time of the first diagnosis of AR. Children who visited the hospital because of some symptoms such as frequent sneezing, rhinorrhea, nasal blockage, and nasal itching, then were diagnosed definitively with AR by the doctor.
- (2) Passive smoking: refers to the exposure of a non-smoker to the smoke emitted from the ignition cigarette or to the smoke exhaled by the smoker for at least 1 day per week.⁷

Statistical analysis

Epi Data 3.1 was used to establish the database, and SPSS 26.0 software was used for data analysis and processing.

Univariate analysis used the χ^2 test. Factors associated with the onset of AR in preschool children were analyzed using multivariate logistic regression, reporting odds ratios (OR), and 95% confidence intervals (CI). All statistical analyses were performed using $\alpha = 0.05$ two-sided test criterion, and $P < 0.05$ was considered statistically significant.

Results

General condition of the investigated subjects

A total of 404 cases were collected in this study, including 225 boys and 179 girls with a mean age of 5.18 ± 0.99 years and a mean age at onset of 3.29 ± 1.11 years. A total of 1616 individuals were matched in the control group in a 1:4 ratio of cases to controls. The control group used both age and sex as matching specifications. There were 900 boys and 716 girls with a mean age 5.13 ± 1.02 years in the control group. The distribution of mode of birth, father with AR, mother with AR, and mother with asthma showed significant differences between the two groups (all $P < 0.05$). There were no significant differences in the distribution of

ethnicity, gestational cycle, birth weight, or duration of breastfeeding between the two groups (all $P > 0.05$) (Table 1).

Univariate analysis of AR and indoor environmental factors

The results of the univariate analysis suggested that renovation or appearance of dampness or mold in the parents' residence before the mother's pregnancy, newly purchased furniture or passive smoking in the parents' residence during the mother's pregnancy, renovation or passive smoking in the children's residence during the child's 0-1 years of age, and father smoking were the main environmental factors contributing to the development of AR in children (all $P < 0.05$) (Table 2).

Multivariate logistic regression analysis of AR onset

General condition survey analysis was introduced into the logistic regression model along with variables that were meaningful in the results of univariate analysis; suffering from AR was the dependent variable (0 = no, 1 = yes). Five

Table 1 Demographics and characteristics of the AR and controls groups.

Factors	AR group (%)	Control group (%)	χ^2	P value
Nation			1.22	0.27
Han	336 (90.6)	1433 (88.7)		
Other ethnic groups	38 (9.4)	183 (11.3)		
Mode of birth			11.45	<0.01
Antepartum	178 (44.1)	864 (53.5)		
Cesarean section	226 (55.9)	752 (46.5)		
Gestational cycles			1.35	0.25
Premature delivery	3 (0.7)	24 (1.5)		
Term birth	401 (99.3)	1592 (98.5)		
Birth weight (g)			1.81	0.41
<2500	15 (3.7)	81 (5.0)		
2500~8000	348 (86.1)	1393 (86.2)		
≥8000	41 (10.1)	142 (8.8)		
Duration of breastfeeding			5.35	0.38
No breastfeeding	23 (5.7)	108 (6.7)		
<1 month	29 (7.2)	85 (5.3)		
1 month - 4 months	48 (11.9)	167 (10.3)		
4 months - 6 months	47 (11.6)	172 (10.6)		
6 months - 1 year	107 (26.5)	492 (30.4)		
≥1 year	150 (37.1)	592 (36.6)		
Father with AR			95.71	<0.01
No	214 (53.0)	1249 (77.3)		
Yes	190 (47.0)	367 (22.7)		
Mother with AR			162.24	<0.01
No	197 (48.8)	1292 (80.0)		
Yes	207 (51.2)	324 (20.0)		
Father with asthma			2.84	0.09
No	397 (98.3)	1603 (99.2)		
Yes	7 (1.7)	13 (0.8)		
Mother with asthma			13.91	<0.01
No	393 (97.3)	1606 (99.4)		
Yes	11 (2.7)	10 (0.6)		

Table 2 Univariate analysis of AR and indoor environmental factors in preschool children between AR and control groups.

Factors	AR group (%)	Control group (%)	χ^2	P value
Furniture was newly purchased in parents' residence before pregnancy			1.85	0.17
No	298 (73.8)	1244 (77.0)		
Yes	106 (26.2)	372 (23.0)		
Renovation in parents' residence before pregnancy			10.02	<0.01
No	319 (79.0)	1380 (85.4)		
Yes	85 (21.0)	236 (14.6)		
Dampness or mold was noted in the parents' residence before pregnancy			5.31	0.02
No	348 (86.1)	1456 (90.1)		
Yes	56 (13.9)	160 (9.9)		
Furniture was newly purchased in the parents' residence during pregnancy			12.86	<0.01
No	338 (83.7)	1454 (90.0)		
Yes	66 (16.3)	162 (10.0)		
Renovation in parents' residence during pregnancy			4.73	0.30
No	367 (90.8)	1517 (93.9)		
Yes	37 (9.2)	99 (6.1)		
Dampness or mold was noted in the parents' residence during pregnancy			3.79	0.05
No	363 (89.9)	1499 (92.8)		
Yes	41 (10.1)	117 (7.2)		
Furniture was newly purchased in children's homes between 0 and 1 year of age			2.13	0.15
No	359 (88.9)	1474 (91.2)		
Yes	45 (11.1)	142 (8.8)		
Residence renovation in children's homes between 0 and 1 year of age			5.23	0.02
No	371 (91.8)	1532 (94.8)		
Yes	33 (8.2)	84 (5.2)		
Dampness or mold was noted in the parents' residence between 0 and 1 year of age			3.82	0.05
No	368 (91.1)	1516 (93.8)		
Yes	36 (8.9)	100 (6.2)		
A hairy pet was kept in children's homes between 0 and 1 year of age			0.00	0.95
No	392 (97.0)	1569 (97.1)		
Yes	12 (3.0)	47 (2.9)		
Flowering plants were grown in children's homes between 0 and 1 year of age			1.04	0.31
No	300 (74.3)	1239 (76.7)		
Yes	104 (25.7)	377 (23.3)		
Father smoking			7.53	<0.01
No	284 (70.3)	1242 (76.9)		
Yes	120 (29.7)	374 (23.1)		
Mother smoking			0.63	0.43
No	403 (99.8)	1607 (99.4)		
Yes	1 (0.2)	9 (0.6)		
Passive smoking in the mother's residence during pregnancy			16.09	<0.01
No	272 (67.3)	1244 (77.0)		
Yes	132 (32.7)	372 (23.0)		
Passive smoking in children's homes between 0 and 1 year of age			15.17	<0.01
No	252 (62.4)	1168 (72.3)		
Yes	152 (37.6)	448 (27.7)		

variables of the final residency main effect model were associated with the onset of AR in preschool children. Mode of birth (cesarean section), father with AR, mother with AR, mother with asthma, and newly purchased furniture in the parents' residence during pregnancy were risk factors for childhood AR (all $P < 0.05$) (Table 3).

Discussion

Previous investigators have suggested that the prevalence of AR is influenced by both genetic⁸ and environmental⁹ factors. This study found that the parental origin of AR contributed significantly to the pathogenesis of AR in children. Concomitant with the ongoing development of genome-wide association studies (GWAS), multiple genes implicated in the pathogenesis of AR have been identified thus far. Within the past 10 years alone, there have been more than 20 investigative studies on genetic susceptibility genes for AR.⁸ A 2018 study by Waage et al.¹⁰ also identified 41 risk loci for AR, which also suggested that AR has high genetic susceptibility. In addition to AR in parents, this investigation identified maternal asthma as a risk factor for AR in children, the mechanism of which is not fully defined. However, this conclusion was similarly reached in a 2007-2008 survey by Wu et al.¹¹

Asthma and AR are both more common allergic diseases with similar aetiologies, often coexisting in the same family or patient.¹² In fact, AR as a significant risk factor for asthma in adults² has been previously reported. As early as 2008, a study from Western Europe¹³ was published in *The Lancet* proposing that either allergic or nonallergic rhinitis could predict the onset of asthma in adults. Regarding how maternal asthma contributes to the onset of AR in children, the authors speculate that on the one hand, there may be an overlap with susceptibility genes. In 2007, the first GWAS on asthma in *Nature* identified a locus (ORMDL3, GSDML, ZPBP2, and IKZF3) on chromosome 17q21 that confers an increased risk of childhood asthma.¹⁴ A subsequent study from Japan¹⁵ also confirmed that variants in this locus were involved in the pathogenesis of AR. On the other hand, AR shares the same environmental influences as asthma. A study by Zhao and colleagues¹⁶ found that passive smoking and petkeeping could increase the risk of AR and asthma in children. In addition, there are many genes associated with asthma and AR pathogenesis and environmental influencing factors that may be responsible for the increased risk of developing AR in children with maternal asthma.

The present investigation found that cesarean section is a risk factor for AR in children. Previous studies have also

suggested that children born by cesarean section have a higher risk of respiratory and allergic diseases than those born vaginally.^{17,18} In particular, elective cesarean section is more likely to cause respiratory morbidity in children.¹⁹ Many studies have confirmed that cesarean section has an important effect on the development and maturation of the neonatal immune system. This may be because children delivered via the birth canal can be exposed to normal beneficial gut microbes from the maternal vagina during delivery. Cesarean-section children are not exposed to these microorganisms, which leads to an imbalance of intestinal flora and the occurrence of diseases such as allergies and autoimmunity.^{20,21} The study by Azad et al.²² also confirmed that infants born by cesarean section had significantly lower richness and diversity of gut flora than those born vaginally, which may be one of the reasons why children born by cesarean section are more susceptible to AR.

For indoor environmental factors, this study only found an association between newly purchased furniture in the parents' residence during the mother's pregnancy and the onset of AR in children. Newly produced furniture, renovation materials, etc., contain large amounts of volatile organic compounds, of which formaldehyde, benzene, toluene, and xylene are the most common substances.²³ Inhalation of such substances at high concentrations irritates the respiratory tract, inducing respiratory disease in children, and formaldehyde is the most representative of these. Many current studies²⁴ have confirmed that formaldehyde can cause respiratory injury. In addition, substances such as formaldehyde can also cross the placental barrier and act on the embryo, causing abnormal fetal growth and development.

Several studies have previously confirmed that indoor passive smoking,¹⁶ parental smoking,²⁵⁻²⁷ keeping houseplants,²⁸ petkeeping,¹⁶ and indoor dampness phenomena²⁹ are closely related to the development of respiratory diseases in children, although this study did not draw the above conclusions. This may be because this study employed a case-control design, and for the case group, subjects were selected strictly according to the physician-diagnosed AR. Some children who presented with AR symptoms but were not definitively diagnosed with AR were not included in the case group. This may have been overly conservative, resulting in the above factors not being found to be associated with AR pathogenesis in children. Additionally, it has been suggested that outdoor air pollution may negatively affect respiratory health but may also enhance sensitization to some plants and fungi.³⁰ In 2012, a "coal to gas" project was launched in Urumqi City, Xinjiang, which greatly improved

Table 3 Multivariate logistic regression analysis of the onset of AR in preschool children between AR and control groups.

Factors	Reference group	β	S-x	Wald χ^2	P value	OR	95%CI
Mode of birth	Antepartum	0.26	0.12	4.49	0.03	1.30	1.02~1.66
Father with AR	No	0.99	0.13	60.74	<0.01	2.68	2.09~3.44
Mother with AR	No	1.32	0.13	108.87	<0.01	3.74	0.92~4.79
Mother with asthma	No	1.09	0.49	4.96	0.03	2.97	1.14~7.76
Furniture was newly purchased in the parents' residence during pregnancy	No	0.58	0.22	6.99	<0.01	1.79	1.16~2.76

the outdoor air quality and may have impacted some factors in the analysis. Therefore, based on the results of the present investigation, it is not yet possible to demonstrate, that the abovementioned factors are not associated with the onset of AR in children. Further confirmation through population-based cohort studies is required to further clarify the contributing factors to the onset of AR in children.

Conclusion

Cesarean section, parents with AR, mother with asthma, and buying new furniture in the mother's home during pregnancy are risk factors for AR in preschool children. The authors should actively carry out AR prevention for cesarean section preschool children with a family history of AR and asthma, and prevent children from contacting harmful factors in the indoor environment early in life.

Data availability

The datasets generated and/or analyzed during the current study are not publicly available due to the data belonging to the School of Public Health of Fudan University but are available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

The study was conducted strictly in accordance with the Declaration of Helsinki and approved by the research ethics committee of Fudan University (protocol no. IRB00002408 & FWA00002399), all parents and class teachers of the children under investigation have signed written informed consent.

Funding

This study was supported by the National Natural Science Foundation of China (grant no. 81861138005, 81860179); Teacher's professional development project of the Shanghai Municipal Education Commission and High-Level Local University Construction Projects of Shanghai University of Medicine & Health Sciences.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Seidman MD, Gurgel RK, Lin SY, Schwartz SR, Baroody FM, Bonner JR, et al. Clinical practice guideline: allergic rhinitis executive summary. *Otolaryngol Head Neck Surg.* 2015;152:197–206.
- Huang K, Yang T, Xu J, Yang L, Zhao J, Zhang X, et al. Prevalence, risk factors, and management of asthma in China: a national cross-sectional study. *Lancet.* 2019;394:407–18.
- Nathan RA. The burden of allergic rhinitis. *Allergy Asthma Proc.* 2007;28:3–9.
- Blaiss MS. Pediatric allergic rhinitis: physical and mental complications. *Allergy Asthma Proc.* 2008;29:1–6.
- Zhang YP, Li BZ, Huang C, Yang X, Qian H, Deng QH, et al. Ten cities cross-sectional questionnaire survey of children asthma and other allergies in China. *Chin Sci Bull.* 2013;58:4182–9.
- To T, Zhu J, Stieb D, Gray N, Fong I, Pinault L, et al. Early life exposure to air pollution and incidence of childhood asthma, allergic rhinitis and eczema. *Eur Respir J.* 2020;55:1900913.
- Luo LF, Chen JW, Luo MH, He ZJ. Exposure to second-hand smoke and its influencing factors among 15-69 years old residents in Guangzhou city. *Chin J Publ Health.* 2019;35:973–6.
- Li J, Zhang Y, Zhang L. Discovering susceptibility genes for allergic rhinitis and allergy using a genome-wide association study strategy. *Curr Opin Allergy Clin Immunol.* 2015;15:33–40.
- Dunlop J, Matsui E, Sharma HP. Allergic rhinitis: environmental determinants. *Immunol Allergy Clin North Am.* 2016;36:367–77.
- Waage J, Standl M, Curtin JA, Jessen LE, Thorsen J, Tian C, et al. Genome-wide association and HLA fine-mapping studies identify risk loci and genetic pathways underlying allergic rhinitis. *Nat Genet.* 2018;50:1072–80. Erratum in: *Nat Genet.* 2018;50:1343.
- Wu KY, Li ZQ, Xue F, Chen W, Jiang MJ, Wang BL, et al. Allergic rhinitis in women and the impact on incidence of allergic diseases in their offspring. *Chin J Otorhinolaryngol-Skull Base Surg.* 2010;16:261–5.
- Rhodes HL, Thomas P, Sporik R, Holgate ST, Cogswell JJ. A birth cohort study of subjects at risk of atopy: twenty-two-year follow-up of wheeze and atopic status. *Am J Respir Crit Care Med.* 2002;165:176–80.
- Shaaban R, Zureik M, Soussan D, Neukirch C, Heinrich J, Sunyer J, et al. Rhinitis and onset of asthma: a longitudinal population-based study. *Lancet.* 2008;372:1049–57.
- Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature.* 2007;448:470–3.
- Tomita K, Sakashita M, Hirota T, Tanaka S, Masuyama K, Yamada T, et al. Variants in the 17q21 asthma susceptibility locus are associated with allergic rhinitis in the Japanese population. *Allergy.* 2013;68:92–100.
- Zhao Y, Liu YQ, Liu MM, Wang D, Ren WH, Gao F, Dong GH. Interactive effects of environmental tobacco smoke and pets ownership on respiratory diseases and symptoms in children. *Zhonghua Er Ke Za Zhi.* 2013;51:96–100. Chinese.
- Hansen AK, Wisborg K, Ulbjerg N, Henriksen TB. Elective caesarean section and respiratory morbidity in the term and near-term neonate. *Acta Obstet Gynecol Scand.* 2007;86:389–94.
- Li SY, Song YJ, Dong JY, Zhao SX, Zhang YW, Jiang YR, et al. Influence of delivery mode on cognition and body development of infants. *Chin J Woman Child Health Res.* 2014;25:922–4.
- Hansen AK, Wisborg K, Ulbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ.* 2008;336:85–7.
- Biasucci G, Benenati B, Morelli L, Bessi E, Boehm G. Cesarean delivery may affect the early biodiversity of intestinal bacteria. *J Nutr.* 2008;138:1796S–800S.
- Renz-Polster H, David MR, Buist AS, Vollmer WM, O'Connor EA, Frazier EA, et al. Cesarean section delivery and the risk of allergic disorders in childhood. *Clin Exp Allergy.* 2005;35:1466–72.
- Azad MB, Konya T, Maughan H, Guttman DS, Field CJ, Chari RS, et al. Gut microbiota of healthy Canadian infants: profiles by mode of delivery and infant diet at 4 months. *CMAJ.* 2013;185:385–94.
- Tang X, Bai Y, Duong A, Smith MT, Li L, Zhang L. Formaldehyde in China: production, consumption, exposure levels, and health effects. *Environ Int.* 2009;35:1210–24.

24. Yan CH, Yi JH. Study progress on mechanism of formaldehyde induced asthma. *China Occupation Med.* 2015;42:682–4.
25. Liu W, Huang C, Hu Y, Zou ZJ, Sundell J. Associations between indoor environmental smoke and respiratory symptoms among preschool children in Shanghai, China. *Chin Sci Bull.* 2013;58:4211–6.
26. Chong-Neto HJ, D'amato G, Rosário Filho NA. Impact of the environment on the microbiome. *J Pediatr.* 2022;98(1):S32–7. Suppl.
27. Ferreira WF, Carvalho DS, Wandalsen GF, Solé D, Sarinho ES, Medeiros D, et al. Associated factors with recurrent wheezing in infants: is there difference between the sexes? *J Pediatr.* 2021;97:629–36.
28. Xie LF, Zhu L, Zhang FZ, Hu WN, Zhao K, Jiang K, Yu W, Sun ZW. Relationship between full flowering stages of anemophilous and ligneous plants and consultation rate of allergic rhinitis. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2016;30:1206–9. Chinese.
29. Zhao ZH, Zhang X, Liu RR, Norback D, Wieslander G, Chen J, et al. Prenatal and early life home environment exposure in relation to preschool children's asthma, allergic rhinitis and eczema in Taiyuan, China. *Chin Sci Bull.* 2013;58:4245–51.
30. Eguluz-Gracia I, Mathioudakis AG, Bartel S, Vijverberg SJ, Fuertes E, Comberiati P, et al. The need for clean air: the way air pollution and climate change affect allergic rhinitis and asthma. *Allergy.* 2020;75:2170–84.