



EDITORIAL

Head growth trajectories as a window into neurodevelopment in preterm infants[☆]

Thiviya Selvanathan  ^{a,b}

^a Department of Pediatrics, University of British Columbia and BC Children's Hospital Research Institute, Vancouver, Canada

^b Department of Pediatrics, The Hospital for Sick Children, Toronto, Canada

Although there have been significant advances in neonatal intensive care over the last few decades, preterm infants remain at increased risk for neurodevelopmental problems that persist into adulthood.¹⁻³ Children born preterm have altered brain maturation, or brain dysmaturation, which includes smaller total and regional brain volumes, altered white matter microstructural maturation, and impaired brain connectivity.⁴ For example, in a study of children born preterm followed to adolescence, slower brain growth and smaller brain volumes were observed in children born preterm compared to children born at term.⁵ Brain dysmaturation in children born preterm has been associated with neurodevelopmental impairments.⁴

Head circumference measurements may provide a window into assessing brain maturation in preterm infants in the clinical setting. Previous studies have shown that head circumference measurements are related to total cerebral volumes in preterm infants.^{6,7} In a cohort of preterm infants with neurodevelopmental follow-up to school-age, a small head circumference at birth was associated with poorer neurodevelopmental outcomes at school-age. However, catch-up growth with normalization of small birth head circumference prior to discharge from the NICU was associated with better outcomes.⁶ Other studies have also observed associations between postnatal head circumference catch-up growth and improved neurodevelopmental outcomes in children born preterm, although findings have varied across cohorts.⁸⁻¹¹

Mayrink et al.¹² followed a prospective cohort of very preterm infants without a history of major neonatal critical illness by performing serial head circumference measurements and neurodevelopmental assessments from birth to

two years of age. There was an overall decrease in head circumference growth during the period of neonatal intensive care with a decrease in z scores from birth to discharge. After discharge from hospital, there was a period of accelerated head growth to 1 month corrected age following which head circumference z scores remained stable. Interestingly, with serial neurodevelopmental assessments performed at 12, 18 and 24 months, there was an increase in neurodevelopmental delays detected in this cohort across timepoints. Larger head circumference at 5 months was associated with higher motor, cognitive and language scores at 18 months corrected age.

The findings of Mayrink et al.¹² reinforce the importance of monitoring head circumference growth in children born preterm. Head circumference measurements can be performed easily in any clinical setting, including those with limited resources and limited access to neuroimaging. Monitoring head circumference trajectories could allow for earlier identification of children who are at increased risk for neurodevelopmental impairments, which may result in implementation of early neurodevelopmental supports and interventions. Specifically, their findings, and those of others, suggest that failure of head circumference catch-up growth in early infancy may be potential marker of an increased risk for neurodevelopmental concerns, although optimal timing of measurements remains unknown.^{6,8,10-12}

The increased prevalence of neurodevelopmental concerns observed over time in the study by Mayrink et al.¹² highlight the importance of longitudinal follow-up to assess neurodevelopmental trajectories in children born preterm. They found an increased prevalence of neurodevelopmental impairments at 24 months compared to 18 months corrected age in their cohort suggesting that later age of assessment may be important for identifying more subtle neurodevelopmental concerns.

[☆] See paper by Mayrink et al. in pages 483–490.

E-mail: thiviya.selvanathan@cw.bc.ca

Their findings are similar to those in a large cohort of very preterm infants with higher odds of a diagnosis of significant developmental impairments in infants who were assessed at 21–24 months compared to children who were assessed at 18–20 months corrected age.¹³ Further, although the authors followed their cohort to 2 years of age, studies examining associations with long-term neurodevelopmental outcomes into childhood and adolescence are important as more subtle impairments may become apparent as children become older.¹⁴ Future studies should also consider assessing neurodevelopmental *trajectories* over time rather than outcomes at a single timepoint.¹⁵ Finally, although the authors excluded preterm infants with major neonatal critical illness in their study and appropriately addressed this as a limitation of their work, future studies of associations between head growth and neurodevelopmental trajectories should also include infants born preterm with major neonatal morbidities so that findings are generalizable to clinical populations of preterm infants.

Conflicts of interest

The author declares no conflicts of interest.

References

- Cheong JL, Olsen JE, Lee KJ, Spittle AJ, Opie GF, Clark M, et al. Temporal trends in neurodevelopmental outcomes to 2 years after extremely preterm birth. *JAMA Pediatr.* 2021;175:1035–42. Erratum in: *JAMA Pediatr.* 2023;177:1242.
- Bell EF, Hintz SR, Hansen NI, Bann CM, Wyckoff MH, DeMauro SB, et al. Mortality, in-hospital morbidity, care practices, and 2-year outcomes for extremely preterm infants in the US, 2013-2018. *JAMA.* 2022;327:248–63. Erratum in: *JAMA.* 2022;327:2151.
- Eves R, Mendonça M, Baumann N, Ni Y, Darlow BA, Horwood J, et al. Association of very preterm birth or very low birth weight with intelligence in adulthood: an individual participant data meta-analysis. *JAMA Pediatr.* 2021;175:e211058.
- Inder TE, Volpe JJ, Anderson PJ. Defining the neurologic consequences of preterm birth. *N Engl J Med.* 2023;389:441–53.
- Thompson DK, Matthews LG, Alexander B, Lee KJ, Kelly CE, Adamson CL, et al. Tracking regional brain growth up to age 13 in children born term and very preterm. *Nat Commun.* 2020;11:696.
- Selvanathan T, Guo T, Kwan E, Chau V, Brant R, Synnes AR, et al. Head circumference, total cerebral volume and neurodevelopment in preterm neonates. *Arch Dis Child Fetal Neonatal Ed.* 2022;107:181–7.
- Cheong JL, Hunt RW, Anderson PJ, Howard K, Thompson DK, Wang HX, et al. Head growth in preterm infants: correlation with magnetic resonance imaging and neurodevelopmental outcome. *Pediatrics.* 2008;121:e1534–40.
- Strobel KM, Wood TR, Valentine GC, German KR, Gogcu S, Hendrixson DT, et al. Contemporary definitions of infant growth failure and neurodevelopmental and behavioral outcomes in extremely premature infants at two years of age. *J Perinatol.* 2024;44:811–8.
- Bando N, Fenton TR, Yang J, Ly L, Luu TM, Unger S, et al. Association of postnatal growth changes and neurodevelopmental outcomes in preterm neonates of <29 weeks' gestation. *J Pediatr.* 2023;256:63–9. e2.
- Sicard M, Nusinovici S, Hanf M, Muller JB, Guellec I, Ancel PY, et al. Fetal and postnatal head circumference growth: synergistic factors for neurodevelopmental outcome at 2 years of age for preterm infants. *Neonatology.* 2017;112:122–9.
- Neubauer V, Griesmaier E, Pehböck-Walser N, Pupp-Peglow U, Kiechl-Kohlendorfer U. Poor postnatal head growth in very preterm infants is associated with impaired neurodevelopment outcome. *Acta Paediatr.* 2013;102:883–8.
- Mayrink ML, Villela LD, Méio MD, Soares FV, de Abranches AD, Nehab SR, et al. The trajectory of head circumference and neurodevelopment in very preterm newborns during the first two years of life: a cohort study. *J Pediatr (Rio J).* 2024;100:483–90.
- Garfinkle J, Khairy M, Simard MN, Wong J, Shah PS, Luu TM, et al. Corrected age at bayley assessment and developmental delay in extreme preterms. *Pediatrics.* 2024;153:e2023063654.
- Christensen R, Chau V, Synnes A, Guo T, Grunau RE, Miller SP. Preterm neurodevelopmental trajectories from 18 months to 4.5 years. *J Pediatr.* 2023;258:113401.
- Neel ML, Conroy S, Srinivas R, Taylor HG, Stark AR, de Silva A, et al. Bayley trajectories predict school readiness better than single assessments in formerly very preterm preschoolers. *Pediatr Res.* 2023;94:1392–9.